


CDC NeVDRS - 2020 and 2021

Summary of Female Suicide Deaths in Nebraska


104 Total Deaths (5.4 per 100,000 Population)








In 2020 and 2021, white females with college credits or Associate's Degree between the age 40-49 most frequently died from suicide. American Indian/Alaska Native NH and Asian/Pacific Islander NH females were disproportionately impacted by suicide in Nebraska.



In 2020 and 2021, approximately 39.5% of female suicides in Nebraska took place between July and October. The month of October recorded the highest number of suicides, accounting for 12.5% of the total. The majority of individuals who lost their lives to suicide during these months were white females, with an average age of 46 and possessing some college credits or Associate's Degree.







Ot	her Dec	edent Characteristics
Statistics are	based on 96 cases	. Categories are not mutually exclusive.
	82.3%	of total suicide deaths took place in a residence.
	10.4%	of total suicide victims experienced homelessness or housing instability.
	63.5%	of total suicide victims had previous mental health problems.
	49.0%	of total suicide victims were diagnosed with depression.
	15.6%	of total suicide victims were diagnosed with bipolar disorder.
	42.7%	of total suicide victims had history of mental health treatment.
	30.2%	of total suicide victims had current mental health treatment.
	59.4%	of total suicide victims were perceived by others to be depressed at the time of death.
	40.6%	of total suicide victims had a history of suicidal ideation.
	26.0%	of total suicide victims had a history of suicide attempt.
	25.0%	of total suicide victims had recently disclosed their suicidal intent.
	11.5%	of total suicide victims had disclosed their suicidal intent to their family member.
	41.7%	of total suicide victims left suicide notes.
	14.6%	of total suicide victims had a history of alcohol abuse.
	33.3%	of total suicide victims had a history of substance abuse other than alcohol.
	20.8%	of total suicide victims had a history of physical health problems.

19.8%

10.4%

36.5%

27.1%

problems.

relationship problems.

married individuals.

individuals.

of total suicide victims had intimate partner

of total suicide victims had family or other

of total suicide victims had known recent

of the suicide deaths were among never

of the suicide deaths were among married