

| Indicators | Overall | | | | | | Men | | | | | | Women | | | | | | LHD Gender Diff. ^e | | | |
|--|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|-------------------------------|------------------------|------------------------------------|------------------|
| | Four Corners | | | State of NE | | | Four Corners | | | State of NE | | | Four Corners | | | State of NE | | | | | | |
| | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d |
| Up-to-date on colon cancer screening, 50-75 year olds^a | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | 301 | 61.5% | (54.8 - 67.7) | 9,022 | 61.1% | (59.6 - 62.5) | NS | 127 | 60.7% | (50.4 - 70.0) | 3,733 | 58.4% | (56.1 - 60.6) | NS | 174 | 62.2% | (53.5 - 70.3) | 5,289 | 63.6% | (61.7 - 65.4) | NS | No |
| 2013 | 293 | 61.1% | (54.6 - 67.3) | 7,979 | 62.8% | (61.1 - 64.4) | NS | 124 | 64.4% | (54.4 - 73.3) | 3,275 | 61.0% | (58.5 - 63.6) | NS | 169 | 57.5% | (48.9 - 65.6) | 4,704 | 64.4% | (62.2 - 66.6) | NS | No |
| 2014 | 425 | 64.9% | (59.4 - 70.0) | 10,646 | 64.1% | (62.8 - 65.5) | NS | 195 | 66.9% | (58.9 - 74.1) | 4,681 | 63.0% | (61.0 - 65.1) | NS | 230 | 62.5% | (55.0 - 69.5) | 5,965 | 65.2% | (63.4 - 67.0) | NS | No |
| 2015 | 291 | 62.1% | (55.6 - 68.2) | 8,007 | 65.2% | (63.6 - 66.7) | NS | 133 | 61.0% | (51.2 - 70.0) | 3,446 | 63.7% | (61.3 - 66.1) | NS | 158 | 63.2% | (54.5 - 71.1) | 4,561 | 66.6% | (64.6 - 68.5) | NS | No |
| 2016 | 312 | 65.0% | (58.7 - 70.8) | 7,036 | 66.0% | (64.3 - 67.6) | NS | 137 | 65.9% | (56.8 - 74.0) | 3,073 | 65.2% | (62.8 - 67.6) | NS | 175 | 63.9% | (55.4 - 71.6) | 3,963 | 66.7% | (64.5 - 68.9) | NS | No |
| Up-to-date on breast cancer screening, female 50-74 year olds^a | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | 173 | 74.4% | (65.9 - 81.4) | 5,200 | 74.9% | (73.2 - 76.5) | NS | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2014 | 226 | 74.8% | (67.9 - 80.6) | 5,904 | 76.1% | (74.5 - 77.7) | NS | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2016 | 170 | 74.9% | (66.9 - 81.4) | 3,914 | 73.4% | (71.3 - 75.4) | NS | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Up-to-date on cervical cancer screening, female 21-65 year olds^a | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | 157 | 82.8% | (73.9 - 89.2) | 5,055 | 83.9% | (82.5 - 85.2) | NS | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2014 | 211 | 76.3% | (68.4 - 82.7) | 5,779 | 81.7% | (80.0 - 83.3) | NS | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2016 | 138 | 79.5% | (70.6 - 86.2) | 3,814 | 77.7% | (75.5 - 79.8) | NS | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Arthritis | | | | | | | | | | | | | | | | | | | | | | |
| Ever told they have arthritis | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 876 | 26.0% | (22.9 - 29.4) | 25,285 | 23.4% | (22.8 - 24.1) | NS | 329 | 21.5% | (17.2 - 26.6) | 10,184 | 20.1% | (19.2 - 21.1) | NS | 547 | 30.5% | (26.3 - 35.0) | 15,101 | 26.6% | (25.7 - 27.6) | NS | No |
| 2012 | 604 | 28.6% | (24.6 - 32.9) | 19,089 | 24.6% | (23.8 - 25.4) | NS | 252 | 26.3% | (20.7 - 32.7) | 7,839 | 21.5% | (20.3 - 22.6) | NS | 352 | 30.8% | (25.4 - 36.8) | 11,250 | 27.7% | (26.6 - 28.8) | NS | No |
| 2013 | 595 | 27.5% | (23.6 - 31.7) | 17,017 | 24.7% | (23.8 - 25.7) | NS | 240 | 25.8% | (20.3 - 32.1) | 6,925 | 21.6% | (20.2 - 23.0) | NS | 355 | 29.2% | (23.9 - 35.1) | 10,092 | 27.8% | (26.5 - 29.2) | NS | No |
| 2014 | 820 | 24.7% | (21.6 - 28.1) | 22,307 | 24.6% | (23.8 - 25.4) | NS | 344 | 19.6% | (15.6 - 24.5) | 9,560 | 21.6% | (20.5 - 22.8) | NS | 476 | 29.7% | (25.2 - 34.6) | 12,747 | 27.4% | (26.3 - 28.6) | NS | Yes |
| 2015 | 666 | 24.4% | (21.1 - 28.2) | 17,472 | 23.4% | (22.6 - 24.3) | NS | 304 | 20.7% | (16.3 - 26.0) | 7,540 | 19.7% | (18.5 - 20.9) | NS | 362 | 28.1% | (23.2 - 33.6) | 9,932 | 27.1% | (25.9 - 28.4) | NS | No |
| 2016 | 604 | 27.0% | (23.1 - 31.4) | 15,137 | 24.6% | (23.6 - 25.6) | NS | 257 | 22.7% | (17.7 - 28.8) | 6,622 | 21.7% | (20.3 - 23.2) | NS | 347 | 31.3% | (25.6 - 37.7) | 8,515 | 27.4% | (26.0 - 28.8) | NS | No |
| Currently have activity limitations due to arthritis, among those ever told they have arthritis^a | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 294 | 43.1% | (36.7 - 49.6) | 7,973 | 45.2% | (43.6 - 46.8) | NS | 88 | 38.3% | (27.8 - 50.1) | 2,711 | 42.6% | (40.0 - 45.2) | NS | 206 | 46.5% | (39.0 - 54.1) | 5,262 | 47.1% | (45.1 - 49.2) | NS | No |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | 195 | 40.5% | (32.9 - 48.6) | 5,460 | 42.4% | (40.3 - 44.6) | NS | 74 | 41.8% | (29.8 - 54.8) | 1,861 | 41.0% | (37.4 - 44.6) | NS | 121 | 39.4% | (30.1 - 49.6) | 3,599 | 43.5% | (40.8 - 46.2) | NS | No |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | 187 | 34.7% | (27.6 - 42.6) | 5,146 | 44.0% | (41.9 - 46.1) | NS | 70 | 29.4% | (19.0 - 42.3) | 1,852 | 41.9% | (38.6 - 45.3) | NS | 117 | 38.3% | (29.2 - 48.4) | 3,294 | 45.5% | (42.9 - 48.1) | NS | No |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Asthma | | | | | | | | | | | | | | | | | | | | | | |
| Ever told they have asthma | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 879 | 10.4% | (8.1 - 13.3) | 25,327 | 11.5% | (10.9 - 12.1) | NS | 328 | 9.7% | (6.3 - 14.5) | 10,194 | 10.5% | (9.6 - 11.3) | NS | 551 | 11.2% | (8.3 - 14.9) | 15,133 | 12.4% | (11.6 - 13.3) | NS | No |
| 2012 | 605 | 9.2% | (6.6 - 12.7) | 19,105 | 10.8% | (10.2 - 11.5) | NS | 252 | 9.9% | (6.0 - 15.9) | 7,847 | 9.7% | (8.8 - 10.7) | NS | 353 | 8.5% | (5.4 - 13.0) | 11,258 | 11.9% | (11.1 - 12.9) | NS | No |
| 2013 | 595 | 8.4% | (6.1 - 11.5) | 17,068 | 11.2% | (10.5 - 12.0) | NS | 240 | 6.9% | (3.9 - 11.8) | 6,946 | 9.6% | (8.6 - 10.8) | NS | 355 | 9.9% | (6.7 - 14.5) | 10,122 | 12.8% | (11.7 - 13.9) | NS | No |
| 2014 | 824 | 11.3% | (8.4 - 15.0) | 22,355 | 12.2% | (11.5 - 13.0) | NS | 347 | 11.9% | (7.3 - 18.8) | 9,583 | 10.8% | (9.8 - 11.9) | NS | 477 | 10.7% | (7.8 - 14.5) | 12,772 | 13.6% | (12.6 - 14.7) | NS | No |
| 2015 | 662 | 8.1% | (5.8 - 11.1) | 17,492 | 12.1% | (11.3 - 12.9) | - | 305 | 6.6% | (3.9 - 11.0) | 7,554 | 10.7% | (9.6 - 12.0) | NS | 357 | 9.6% | (6.3 - 14.4) | 9,938 | 13.4% | (12.3 - 14.5) | NS | No |
| 2016 | 605 | 15.4% | (11.4 - 20.6) | 15,144 | 12.4% | (11.5 - 13.4) | NS | 257 | 15.0% | (9.5 - 22.9) | 6,622 | 10.4% | (9.2 - 11.8) | NS | 348 | 15.8% | (10.4 - 23.4) | 8,522 | 14.3% | (13.1 - 15.7) | NS | No |
| Currently have asthma | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 877 | 5.8% | (4.2 - 8.0) | 25,257 | 7.3% | (6.9 - 7.8) | NS | 326 | 4.6% | (2.5 - 8.4) | 10,170 | 6.0% | (5.4 - 6.7) | NS | 551 | 7.0% | (4.9 - 9.9) | 15,087 | 8.6% | (8.0 - 9.4) | NS | No |
| 2012 | 604 | 7.0% | (4.8 - 10.3) | 19,053 | 7.4% | (6.9 - 7.9) | NS | 251 | 7.5% | (4.1 - 13.1) | 7,832 | 6.1% | (5.4 - 6.9) | NS | 353 | 6.6% | (4.0 - 11.0) | 11,221 | 8.7% | (7.9 - 9.4) | NS | No |
| 2013 | 593 | 6.9% | (4.7 - 9.9) | 17,024 | 7.3% | (6.7 - 8.0) | NS | 239 | 6.0% | (3.3 - 11.0) | 6,930 | 5.5% | (4.8 - 6.4) | NS | 354 | 7.7% | (4.8 - 12.1) | 10,094 | 9.1% | (8.2 - 10.0) | NS | No |
| 2014 | 822 | 6.6% | (4.7 - 9.1) | 22,289 | 7.7% | (7.2 - 8.3) | NS | 347 | 5.7% | (3.3 - 9.8) | 9,562 | 6.2% | (5.5 - 7.0) | NS | 475 | 7.4% | (5.0 - 10.8) | 12,727 | 9.2% | (8.4 - 10.1) | NS | No |
| 2015 | 659 | 5.6% | (3.7 - 8.3) | 17,435 | 7.2% | (6.6 - 7.8) | NS | 305 | 4.6% | (2.3 - 9.0) | 7,533 | 5.4% | (4.7 - 6.2) | NS | 354 | 6.5% | (3.9 - 10.6) | 9,902 | 8.9% | (8.0 - 9.9) | NS | No |
| 2016 | 604 | 7.7% | (5.2 - 11.3) | 15,116 | 8.3% | (7.5 - 9.1) | NS | 257 | 6.8% | (3.9 - 11.8) | 6,609 | 6.1% | (5.2 - 7.1) | NS | 347 | 8.6% | (4.9 - 14.5) | 8,507 | 10.4% | (9.3 - 11.6) | NS | No |

| Indicators | Overall | | | | | | Men | | | | | | Women | | | | | | LHD Gender Diff. ^e | | | | |
|--|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|-------------------------------|------------------------|------------------------------------|------------------|---|
| | Four Corners | | | State of NE | | | Four Corners | | | State of NE | | | Four Corners | | | State of NE | | | | | | | |
| | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | |
| Current e-cigarette use | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2016 | 589 | 4.3% | (2.3 - 7.7) | 14,819 | 4.9% | (4.3 - 5.6) | NS | 250 | 4.9% | (2.3 - 10.2) | 6,474 | 5.3% | (4.5 - 6.4) | NS | 339 | 3.7% | (1.4 - 9.4) | 8,345 | 4.5% | (3.7 - 5.6) | NS | No | |
| Overweight and Obesity | | | | | | | | | | | | | | | | | | | | | | | |
| Obese (BMI=30+) [^] | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 846 | 28.3% | (24.6 - 32.2) | 24,366 | 28.4% | (27.6 - 29.2) | NS | 330 | 28.1% | (22.8 - 34.1) | 10,102 | 29.2% | (28.0 - 30.4) | NS | 516 | 28.4% | (23.7 - 33.7) | 14,264 | 27.6% | (26.5 - 28.8) | NS | No | |
| 2012 | 587 | 27.6% | (23.3 - 32.4) | 18,385 | 28.6% | (27.7 - 29.6) | NS | 251 | 30.3% | (23.9 - 37.4) | 7,781 | 29.2% | (27.9 - 30.6) | NS | 336 | 25.0% | (19.4 - 31.5) | 10,604 | 28.1% | (26.8 - 29.3) | NS | No | |
| 2013 | 573 | 32.4% | (27.7 - 37.5) | 16,250 | 29.6% | (28.4 - 30.7) | NS | 240 | 35.2% | (28.0 - 43.1) | 6,860 | 30.8% | (29.1 - 32.5) | NS | 333 | 29.4% | (23.7 - 35.9) | 9,390 | 28.4% | (26.9 - 29.9) | NS | No | |
| 2014 | 788 | 31.2% | (27.3 - 35.3) | 21,130 | 30.2% | (29.2 - 31.3) | NS | 344 | 31.1% | (25.6 - 37.1) | 9,399 | 31.7% | (30.2 - 33.2) | NS | 444 | 31.3% | (26.2 - 36.9) | 11,731 | 28.7% | (27.4 - 30.1) | NS | No | |
| 2015 | 613 | 33.6% | (29.1 - 38.3) | 16,265 | 31.4% | (30.3 - 32.5) | NS | 295 | 32.8% | (26.8 - 39.4) | 7,323 | 32.9% | (31.3 - 34.6) | NS | 318 | 34.4% | (28.0 - 41.5) | 8,942 | 29.8% | (28.3 - 31.3) | NS | No | |
| 2016 | 573 | 29.1% | (24.9 - 33.7) | 14,173 | 32.0% | (30.8 - 33.2) | NS | 253 | 33.3% | (26.8 - 40.4) | 6,460 | 32.6% | (30.8 - 34.3) | NS | 320 | 24.6% | (19.7 - 30.3) | 7,713 | 31.4% | (29.7 - 33.1) | NS | No | |
| Overweight or Obese (BMI=25+) | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 846 | 66.1% | (61.8 - 70.2) | 24,366 | 64.9% | (64.0 - 65.8) | NS | 330 | 70.8% | (63.8 - 76.9) | 10,102 | 72.8% | (71.5 - 74.1) | NS | 516 | 61.1% | (55.7 - 66.3) | 14,264 | 56.8% | (55.5 - 58.1) | NS | No | |
| 2012 | 587 | 63.7% | (58.5 - 68.6) | 18,385 | 65.0% | (64.0 - 66.0) | NS | 251 | 67.3% | (59.4 - 74.4) | 7,781 | 72.3% | (70.8 - 73.7) | NS | 336 | 60.0% | (53.1 - 66.6) | 10,604 | 57.6% | (56.2 - 59.0) | NS | No | |
| 2013 | 573 | 68.9% | (63.5 - 73.8) | 16,250 | 65.5% | (64.2 - 66.7) | NS | 240 | 73.7% | (65.6 - 80.4) | 6,860 | 72.2% | (70.4 - 74.0) | NS | 333 | 63.8% | (56.4 - 70.5) | 9,390 | 58.5% | (56.8 - 60.2) | NS | No | |
| 2014 | 788 | 72.8% | (68.6 - 76.7) | 21,130 | 66.7% | (65.6 - 67.8) | + | 344 | 78.6% | (72.6 - 83.6) | 9,399 | 73.8% | (72.3 - 75.3) | NS | 444 | 66.8% | (60.6 - 72.3) | 11,731 | 59.2% | (57.7 - 60.7) | NS | Yes | |
| 2015 | 613 | 70.1% | (65.4 - 74.5) | 16,265 | 67.0% | (65.8 - 68.2) | NS | 295 | 73.5% | (66.6 - 79.4) | 7,323 | 73.4% | (71.7 - 75.1) | NS | 318 | 66.4% | (59.6 - 72.6) | 8,942 | 60.2% | (58.6 - 61.9) | NS | No | |
| 2016 | 573 | 66.7% | (61.0 - 71.9) | 14,173 | 68.5% | (67.3 - 69.8) | NS | 253 | 79.6% | (72.0 - 85.5) | 6,460 | 74.9% | (73.1 - 76.6) | NS | 320 | 52.8% | (45.3 - 60.3) | 7,713 | 61.8% | (59.9 - 63.6) | NS | Yes | |
| Nutrition | | | | | | | | | | | | | | | | | | | | | | | |
| Consumed sugar-sweetened beverages 1 or more times per day in past 30 days | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | 269 | 26.6% | (20.1 - 34.3) | 7,822 | 28.5% | (26.8 - 30.3) | NS | 106 | 33.2% | (23.1 - 45.3) | 3,164 | 35.8% | (33.1 - 38.5) | NS | 163 | 20.8% | (13.2 - 31.2) | 4,658 | 21.7% | (19.7 - 24.0) | NS | No | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Currently watching or reducing sodium or salt intake | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | 270 | 38.9% | (32.5 - 45.8) | 7,818 | 46.3% | (44.6 - 48.1) | NS | 107 | 40.9% | (30.8 - 51.8) | 3,165 | 43.4% | (40.8 - 46.0) | NS | 163 | 37.2% | (29.1 - 46.0) | 4,653 | 49.1% | (46.8 - 51.5) | - | No | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | 293 | 50.4% | (42.5 - 58.3) | 7,892 | 46.8% | (45.0 - 48.5) | NS | 129 | 44.0% | (33.2 - 55.4) | 3,436 | 46.0% | (43.4 - 48.7) | NS | 164 | 57.3% | (45.4 - 68.3) | 4,456 | 47.5% | (45.2 - 49.8) | NS | No | |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Median times per day consumed fruits | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 836 | 1.02 | (1.00 - 1.15) | 24,090 | 1.02 | (1.02 - 1.04) | NS | 317 | 0.81 | (0.68 - 0.91) | 9,660 | 0.99 | (0.99 - 1.00) | - | 519 | 1.42 | (1.21 - 1.60) | 14,430 | 1.14 | (1.13 - 1.14) | + | Yes | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | 563 | 1.00 | (0.98 - 1.06) | 15,949 | 1.00 | (1.00 - 1.00) | NS | 229 | 0.96 | (0.72 - 0.99) | 6,462 | 0.99 | (0.99 - 1.00) | - | 334 | 1.07 | (1.01 - 1.42) | 9,487 | 1.07 | (1.06 - 1.14) | NS | Yes | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | 609 | 0.99 | (0.96 - 1.07) | 16,135 | 1.00 | (1.00 - 1.00) | NS | 278 | 0.93 | (0.72 - 0.99) | 6,899 | 0.99 | (0.98 - 1.00) | NS | 331 | 1.07 | (0.98 - 1.53) | 9,236 | 1.03 | (1.02 - 1.09) | NS | No | |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Consumed fruits less than 1 time per day | | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 836 | 42.0% | (37.7 - 46.5) | 24,090 | 40.1% | (39.2 - 41.1) | NS | 317 | 55.1% | (48.5 - 61.5) | 9,660 | 46.2% | (44.8 - 47.6) | + | 519 | 28.8% | (23.9 - 34.1) | 14,430 | 34.3% | (33.1 - 35.6) | NS | Yes | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | 563 | 40.3% | (35.0 - 45.9) | 15,949 | 39.7% | (38.5 - 41.0) | NS | 229 | 49.3% | (41.3 - 57.3) | 6,462 | 45.4% | (43.4 - 47.3) | NS | 334 | 31.2% | (24.6 - 38.8) | 9,487 | 34.4% | (32.8 - 36.1) | NS | Yes | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2015 | 609 | 39.1% | (34.3 - 44.1) | 16,135 | 41.1% | (39.9 - 42.4) | NS | 278 | 47.1% | (40.1 - 54.2) | 6,899 | 47.4% | (45.6 - 49.3) | NS | 331 | 31.0% | (24.8 - 38.0) | 9,236 | 35.1% | (33.5 - 36.7) | NS | Yes | |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |

| Indicators | Overall | | | | | | Men | | | | | | Women | | | | | | LHD Gender Diff. ^e | | | |
|--|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|-------------------------------|------------------------|------------------|------------------------------------|
| | Four Corners | | | State of NE | | | Sig ^d | Four Corners | | | State of NE | | | Sig ^d | Four Corners | | | State of NE | | | Sig ^d | |
| | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | | mean or % ^b | | 95% C.I. ^c (Low - High) |
| Immunization and Infectious Disease | | | | | | | | | | | | | | | | | | | | | | |
| Had a flu vaccination in past year, aged 18 years and older | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 854 | 36.7% | (32.9 - 40.7) | 24,156 | 41.1% | (40.2 - 42.0) | NS | 323 | 32.3% | (26.7 - 38.5) | 9,663 | 35.1% | (33.8 - 36.5) | NS | 531 | 41.2% | (36.3 - 46.3) | 14,493 | 46.8% | (45.6 - 48.1) | NS | No |
| 2012 | 600 | 42.2% | (37.4 - 47.2) | 18,903 | 42.2% | (41.2 - 43.2) | NS | 250 | 38.3% | (31.4 - 45.7) | 7,754 | 36.8% | (35.4 - 38.2) | NS | 350 | 46.1% | (39.5 - 52.9) | 11,149 | 47.5% | (46.1 - 48.8) | NS | No |
| 2013 | 554 | 44.9% | (39.6 - 50.4) | 16,022 | 45.2% | (43.9 - 46.4) | NS | 226 | 41.5% | (34.0 - 49.4) | 6,506 | 41.2% | (39.4 - 43.1) | NS | 328 | 48.5% | (41.2 - 55.0) | 9,516 | 48.9% | (47.2 - 50.6) | NS | No |
| 2014 | 803 | 41.0% | (36.9 - 45.3) | 21,638 | 43.9% | (42.9 - 45.0) | NS | 341 | 37.6% | (31.6 - 44.0) | 9,262 | 38.8% | (37.3 - 40.4) | NS | 462 | 44.5% | (38.9 - 50.2) | 12,376 | 48.9% | (47.4 - 50.3) | NS | No |
| 2015 | 606 | 45.7% | (40.9 - 50.6) | 16,091 | 47.2% | (46.0 - 48.4) | NS | 276 | 43.8% | (37.0 - 50.8) | 6,883 | 40.8% | (39.0 - 42.6) | NS | 330 | 47.6% | (40.8 - 54.4) | 9,208 | 53.4% | (51.7 - 55.0) | NS | No |
| 2016 | 582 | 38.9% | (34.1 - 44.0) | 14,685 | 44.4% | (43.1 - 45.7) | NS | 247 | 34.6% | (27.8 - 42.1) | 6,401 | 39.4% | (37.5 - 41.3) | NS | 335 | 43.3% | (36.6 - 50.3) | 8,284 | 49.3% | (47.6 - 51.1) | NS | No |
| Had a flu vaccination in past year, aged 65 years and older ^a | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 312 | 56.6% | (50.3 - 62.6) | 8,185 | 61.8% | (60.3 - 63.3) | NS | 102 | 54.9% | (44.4 - 64.9) | 2,875 | 59.6% | (57.1 - 62.1) | NS | 210 | 57.9% | (50.2 - 65.2) | 5,310 | 63.4% | (61.5 - 65.3) | NS | No |
| 2012 | 216 | 65.8% | (58.2 - 72.7) | 6,625 | 62.9% | (61.2 - 64.6) | NS | 86 | 63.0% | (51.0 - 73.6) | 2,327 | 60.9% | (58.1 - 63.6) | NS | 130 | 68.7% | (59.3 - 76.8) | 4,298 | 64.4% | (62.4 - 66.5) | NS | No |
| 2013 | 199 | 64.2% | (56.5 - 71.2) | 5,652 | 66.2% | (64.2 - 68.1) | NS | 78 | 67.6% | (55.2 - 77.8) | 2,067 | 67.6% | (64.4 - 70.6) | NS | 121 | 61.2% | (51.4 - 70.2) | 3,585 | 65.0% | (62.6 - 67.4) | NS | No |
| 2014 | 297 | 58.8% | (52.1 - 65.2) | 7,610 | 64.7% | (63.2 - 66.2) | NS | 130 | 53.0% | (43.2 - 62.5) | 2,971 | 65.9% | (63.5 - 68.3) | - | 167 | 65.4% | (56.8 - 73.1) | 4,639 | 63.8% | (61.8 - 65.7) | NS | No |
| 2015 | 205 | 62.5% | (54.4 - 69.9) | 5,578 | 65.2% | (63.4 - 67.0) | NS | 87 | 63.3% | (51.2 - 73.9) | 2,119 | 64.1% | (61.1 - 66.9) | NS | 118 | 61.7% | (50.6 - 71.7) | 3,459 | 66.1% | (63.9 - 68.2) | NS | No |
| 2016 | 224 | 60.1% | (52.4 - 67.3) | 5,130 | 62.7% | (60.8 - 64.6) | NS | 83 | 59.4% | (47.1 - 70.6) | 1,967 | 62.8% | (59.8 - 65.6) | NS | 141 | 60.7% | (51.2 - 69.5) | 3,163 | 62.7% | (60.2 - 65.1) | NS | No |
| Ever had a pneumonia vaccination, aged 65 years and older ^a | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | 300 | 64.6% | (58.1 - 70.5) | 7,997 | 70.3% | (68.8 - 71.7) | NS | 96 | 61.2% | (50.3 - 71.0) | 2,772 | 66.7% | (64.2 - 69.1) | NS | 204 | 67.2% | (59.4 - 74.1) | 5,225 | 72.9% | (71.2 - 74.5) | NS | No |
| 2012 | 209 | 68.3% | (60.7 - 75.0) | 6,413 | 70.0% | (68.4 - 71.6) | NS | 82 | 70.4% | (58.7 - 79.8) | 2,225 | 66.9% | (64.1 - 69.6) | NS | 127 | 66.2% | (56.1 - 75.0) | 4,188 | 72.3% | (70.3 - 74.1) | NS | No |
| 2013 | 191 | 68.8% | (61.3 - 75.4) | 5,478 | 71.7% | (69.8 - 73.4) | NS | 75 | 76.2% | (64.5 - 84.9) | 1,970 | 71.5% | (68.5 - 74.3) | NS | 116 | 62.4% | (52.4 - 71.5) | 3,508 | 71.8% | (69.4 - 74.0) | NS | No |
| 2014 | 286 | 68.0% | (61.4 - 73.9) | 7,372 | 72.3% | (70.9 - 73.7) | NS | 123 | 62.8% | (52.9 - 71.7) | 2,849 | 70.3% | (67.9 - 72.5) | NS | 163 | 73.6% | (65.0 - 80.7) | 4,523 | 73.9% | (72.1 - 75.7) | NS | No |
| 2015 | 200 | 72.2% | (64.5 - 78.8) | 5,452 | 73.8% | (72.1 - 75.4) | NS | 86 | 69.8% | (57.9 - 79.5) | 2,056 | 70.3% | (67.4 - 73.0) | NS | 114 | 74.4% | (64.1 - 82.6) | 3,396 | 76.5% | (74.5 - 78.4) | NS | No |
| 2016 | 219 | 74.9% | (68.0 - 80.8) | 5,060 | 75.9% | (74.2 - 77.5) | NS | 81 | 76.3% | (65.4 - 84.6) | 1,927 | 73.7% | (70.9 - 76.3) | NS | 138 | 73.7% | (64.5 - 81.2) | 3,133 | 77.6% | (75.5 - 79.7) | NS | No |
| Had a tetanus vaccination since 2005 | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| 2013 | 509 | 59.6% | (54.0 - 65.0) | 14,650 | 60.2% | (58.9 - 61.5) | NS | 201 | 64.0% | (54.9 - 72.1) | 6,015 | 63.1% | (61.1 - 65.0) | NS | 308 | 55.5% | (48.1 - 62.7) | 8,635 | 57.4% | (55.6 - 59.1) | NS | No |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | | |
| 2016 | 537 | 68.5% | (63.7 - 72.9) | 13,574 | 64.2% | (62.9 - 65.4) | NS | 231 | 71.7% | (64.8 - 77.7) | 5,939 | 67.5% | (65.5 - 69.3) | NS | 306 | 65.2% | (58.3 - 71.5) | 7,635 | 60.9% | (59.2 - 62.7) | NS | No |

| Indicators | Overall | | | | | | Men | | | | | | Women | | | | | | LHD Gender Diff. ^e | | | |
|--|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|-------------------------------|------------------------|------------------|------------------------------------|
| | Four Corners | | | State of NE | | | Sig ^d | Four Corners | | | State of NE | | | Sig ^d | Four Corners | | | State of NE | | | Sig ^d | |
| | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | | mean or % ^b | | 95% C.I. ^c (Low - High) |
| Had teeth cleaned by dentist/hygienist in past year, among those with 1+ permanent teeth | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2013 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2016 | 561 | 67.7% | (62.4 - 72.5) | 13,764 | 67.4% | (66.1 - 68.6) | NS | 239 | 64.3% | (56.2 - 71.6) | 6,092 | 63.9% | (62.0 - 65.8) | NS | 322 | 71.0% | (64.2 - 77.0) | 7,672 | 70.8% | (69.1 - 72.5) | NS | No |
| Social Context | | | | | | | | | | | | | | | | | | | | | | |
| Housing insecurity in past year, among those who own or rent their home [^] | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | 242 | 18.7% | (12.7 - 26.5) | 6,089 | 26.6% | (25.0 - 28.3) | NS | 102 | 18.5% | (10.7 - 30.0) | 2,481 | 24.7% | (22.3 - 27.3) | NS | 140 | 18.8% | (10.9 - 30.6) | 3,608 | 28.5% | (26.3 - 30.8) | NS | No |
| 2013 | 259 | 18.3% | (13.4 - 24.6) | 7,324 | 28.8% | (27.1 - 30.5) | - | 99 | 14.7% | (8.3 - 24.6) | 2,961 | 25.0% | (22.5 - 27.7) | NS | 160 | 21.2% | (14.7 - 29.7) | 4,363 | 32.3% | (30.0 - 34.7) | - | No |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2015 | 293 | 31.4% | (24.9 - 38.9) | 7,461 | 28.5% | (26.8 - 30.2) | NS | 138 | 25.5% | (17.7 - 35.4) | 3,109 | 26.6% | (24.2 - 29.2) | NS | 155 | 37.3% | (27.2 - 48.5) | 4,352 | 30.3% | (28.1 - 32.5) | NS | No |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Food insecurity in past year [^] | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | 263 | 15.8% | (9.8 - 24.5) | 6,621 | 17.6% | (16.2 - 19.0) | NS | 114 | 9.8% | (4.3 - 20.6) | 2,684 | 13.9% | (12.1 - 16.0) | NS | 149 | 21.9% | (12.3 - 35.9) | 3,937 | 21.1% | (19.1 - 23.2) | NS | No |
| 2013 | 269 | 11.4% | (7.5 - 16.8) | 7,828 | 19.0% | (17.6 - 20.6) | - | 106 | 11.8% | (6.1 - 21.6) | 3,159 | 15.4% | (13.4 - 17.7) | NS | 163 | 11.0% | (6.6 - 17.8) | 4,669 | 22.5% | (20.5 - 24.6) | - | No |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2015 | 305 | 23.6% | (17.6 - 31.0) | 7,918 | 21.0% | (19.5 - 22.5) | NS | 145 | 21.3% | (14.0 - 31.2) | 3,309 | 17.9% | (15.8 - 20.2) | NS | 160 | 25.9% | (17.0 - 37.3) | 4,609 | 24.0% | (22.1 - 26.1) | NS | No |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Caregiving | | | | | | | | | | | | | | | | | | | | | | |
| Provided regular care/assistance in past month to friend or family member with health issue | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2013 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2015 | 292 | 25.0% | (18.3 - 33.1) | 7,883 | 24.7% | (23.2 - 26.2) | NS | 130 | 20.1% | (11.5 - 32.7) | 3,434 | 21.1% | (19.1 - 23.4) | NS | 162 | 30.4% | (21.0 - 41.7) | 4,449 | 28.0% | (25.9 - 30.2) | NS | No |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Cognitive Decline | | | | | | | | | | | | | | | | | | | | | | |
| Experienced more or worsening confusion or memory loss in past year, aged 45 years and older | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2013 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2014 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2015 | 215 | 9.1% | (5.5 - 14.7) | 5,767 | 9.4% | (8.4 - 10.5) | NS | 89 | 13.7% | (7.2 - 24.5) | 2,436 | 10.2% | (8.7 - 12.0) | NS | 126 | 5.3% | (2.6 - 10.7) | 3,331 | 8.6% | (7.3 - 10.2) | NS | No |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Inadequate Sleep | | | | | | | | | | | | | | | | | | | | | | |
| Get less than 7 hours of sleep per day | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2013 | 592 | 26.3% | (22.1 - 31.1) | 16,972 | 31.8% | (30.6 - 33.0) | NS | 239 | 27.8% | (21.3 - 35.4) | 6,927 | 32.6% | (30.8 - 34.4) | NS | 353 | 24.9% | (19.8 - 30.8) | 10,045 | 31.0% | (29.5 - 32.6) | NS | No |
| 2014 | 808 | 30.1% | (26.1 - 34.5) | 22,172 | 30.0% | (29.0 - 31.0) | NS | 341 | 30.5% | (24.4 - 37.4) | 9,527 | 30.7% | (29.2 - 32.2) | NS | 467 | 29.7% | (24.7 - 35.3) | 12,645 | 29.4% | (28.1 - 30.7) | NS | No |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2016 | 605 | 26.7% | (22.4 - 31.4) | 15,097 | 29.6% | (28.4 - 30.8) | NS | 256 | 29.0% | (22.4 - 36.6) | 6,610 | 29.7% | (27.9 - 31.5) | NS | 349 | 24.3% | (19.2 - 30.3) | 8,487 | 29.5% | (27.9 - 31.2) | NS | No |
| Average hours of sleep per day | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2013 | 592 | 7.1 | (7.0 - 7.3) | 16,972 | 7.1 | (7.0 - 7.1) | NS | 239 | 7.2 | (7.0 - 7.3) | 6,927 | 7.0 | (7.0 - 7.1) | NS | 353 | 7.1 | (7.0 - 7.3) | 10,045 | 7.1 | (7.0 - 7.1) | NS | No |
| 2014 | 808 | 7.1 | (7.0 - 7.2) | 22,172 | 7.1 | (7.0 - 7.1) | NS | 341 | 7.0 | (6.9 - 7.2) | 9,527 | 7.0 | (7.0 - 7.1) | NS | 467 | 7.1 | (7.0 - 7.2) | 12,645 | 7.1 | (7.1 - 7.1) | NS | No |
| 2015 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| 2016 | 605 | 7.2 | (7.1 - 7.3) | 15,097 | 7.1 | (7.1 - 7.1) | NS | 256 | 7.1 | (7.0 - 7.3) | 6,610 | 7.1 | (7.0 - 7.2) | NS | 349 | 7.3 | (7.0 - 7.5) | 8,487 | 7.1 | (7.1 - 7.2) | NS | No |

| Indicators | Overall | | | | | | Men | | | | | | Women | | | | | | LHD Gender Diff. ^e | | | |
|---|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|------------------------|------------------------------------|------------------|----------------|------------------------|------------------------------------|----------------|-------------------------------|------------------------|------------------------------------|------------------|
| | Four Corners | | | State of NE | | | Four Corners | | | State of NE | | | Four Corners | | | State of NE | | | | | | |
| | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d | n ^a | mean or % ^b | 95% C.I. ^c (Low - High) | n ^a | | mean or % ^b | 95% C.I. ^c (Low - High) | Sig ^d |
| Occupational Safety and Health | | | | | | | | | | | | | | | | | | | | | | |
| Work-related injury or illness in past year, among employed or recently out of work | | | | | | | | | | | | | | | | | | | | | | |
| 2011 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2012 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 2013 | 173 | 5.5% | (2.7 - 10.8) | 4,952 | 4.8% | (3.9 - 5.8) | NS | 82 | 8.3% | (3.8 - 17.4) | 2,246 | 6.1% | (4.8 - 7.7) | NS | 91 | 1.8% | (0.4 - 7.2) | 2,706 | 3.3% | (2.2 - 4.9) | NS | No |
| 2014 | 259 | 1.4% | (0.6 - 3.4) | 6,966 | 4.9% | (4.1 - 5.9) | - | 116 | 1.0% | (0.3 - 3.4) | 3,403 | 5.4% | (4.4 - 6.7) | - | 143 | 1.8% | (0.5 - 5.9) | 3,563 | 4.3% | (3.1 - 6.0) | NS | No |
| 2015 | 203 | 4.7% | (2.2 - 10.0) | 5,434 | 5.7% | (4.7 - 6.8) | NS | 103 | 5.3% | (1.8 - 15.0) | 2,727 | 7.1% | (5.6 - 9.0) | NS | 100 | 3.9% | (1.6 - 8.9) | 2,707 | 3.9% | (2.9 - 5.3) | NS | No |
| 2016 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |

Note: Data reflect the four county LHD region of Butler, Polk, Seward, and York Counties

Note: Data reflect both landline and cell phone responses

Note: This table is not intended to be inclusive of all BRFSS indicators; some were excluded due to small numbers at the LHD level

Note: This table excludes 2011 BRFSS optional module and state added questions data due to the data being landline only

Note: The results in this table may be slightly different than results in previously published reports and tables due differences in the software used and/or how the statistical analysis code was written; these results were analyzed using SAS and SAS-callable SUDAAN software

^a Non-weighted sample size among adults 18 and older (unless different age group noted)

^b Weighted mean, median, or percentage (percentages are followed by the % symbol) among adults 18 and older (unless different age group noted)

^c Low and High are the lower and upper limits of the 95% confidence interval, respectively

^d Symbols represent: "+" = LHD value significantly higher than the state value; "-" = LHD value significantly lower than the state value; "NS" = LHD value not statistically different than the state value; significant differences based on 95% confidence interval overlap

^e Indicates whether there is a significant difference by gender within the LHD region, based on non-overlapping 95% confidence intervals; note that small sample sizes for males and females can make interpreting significance difficult due to large confidence intervals

[^] Reflects a Nebraska Healthy People 2020 (HP2020) measure

* Data suppressed due to a small number of respondents (i.e., fewer than 50)

Source: Nebraska Behavioral Risk Factor Surveillance System (BRFSS); January 2018

