

## CHILDHOOD IMMUNIZATION CHART

Child's Age at Entrance or at Time of Record Review	<<<<Minimum>>>> Number of Doses Required of Each Vaccine						
	PCV7 or 13 Vaccine	DTaP / DT Td / Tdap Vaccine	Polio Vaccine	Hib Vaccine	Hep B Vaccine	MMR MMRV Vaccine	Varicella Vaccine VZV
0-But Not Yet 3 Months	None	None	None	None	None	None	None
3 Months But Not Yet 5 Months	1 Dose	1 Dose	1 Dose	1 Dose	1 Dose	None	None
5 Months But Not Yet 7 Months	2 Doses	2 Doses	2 Doses	2 Doses	2 Doses	None	None
7 Months But Not Yet 16 Months	3 Doses * * * *	3 Doses	2 Doses	2 Doses **	3 Doses	None	None
16 Months But Not Yet 19 Months	4 Doses * * * *	3 Doses	2 Doses	3 Doses **	3 Doses	1 Dose *	1 Dose * or documented history of disease
19 Months to School Entry	4 Doses * * * *	4 Doses	3 Doses	3 Doses **	3 Doses	1 Dose *	1 Dose * or documented history of disease
At School Entry	None	4 Doses	3 Doses	None ***	3 Doses	2 Dose *	2 Dose * or documented history of disease

\* First dose must be given at or after 12 months of age

\*\* Hib vaccine is recommended to be given in a multiple dose schedule beginning in infancy. However, any child who has received a single dose of Hib vaccine at or after 15 months of age is appropriately immunized.

\*\*\* Hib vaccine not required after child reaches 5 years of age.

\*\*\*\* If a child is 7 months of age or older when they start the PCV series, they do not need all 4 doses of PCV. Unvaccinated children who begin the PCV series between 7 – 11 months should receive 2 doses 4 weeks apart with a booster at 12 – 15 months. Unvaccinated children who begin the PCV series between 12 – 23 months of age should receive 2 doses 8 weeks apart. Unvaccinated children 24 – 59 months of age need 1 dose of PCV7. Children with high-risk conditions age 24 – 59 months should follow their physician's advice.