

Table 1C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Demographic Indicators ⁽²⁾

Total Records	Number	%	Source of Data	Number	%
Complete Records	8,566	74.2	WIC	11,549	100.0
Prenatal Records	766	6.6	Maternal Health	0	0.0
Postpartum Records	2,217	19.2	Other	0	0.0
Total Records Accepted	11,549		Total	11,549	100.0
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%
White, Not Hispanic	6,575	56.9
Black, Not Hispanic	1,234	10.7
Hispanic	3,224	27.9
American Indian/Alaska Native	179	1.6
Asian/Pacific Islander	213	1.8
Multiple Races	122	1.1
All Other	0	0.0
Total	11,547	100.0
<i>Excluded Unknown: 2</i>	<i>0.0</i>	

Age Distribution	Number	%
< 15 Years	35	0.3
15 - 17 Years	667	5.8
18 - 19 Years	1,420	12.3
20 - 29 Years	7,208	62.5
30 - 39 Years	2,063	17.9
>= 40 Years	149	1.3
Total	11,542	100.0
<i>Excluded Unknown: 7</i>	<i>0.1</i>	

Education Distribution	Number	%
< High School	3,758	32.6
High School	4,581	39.8
> High School	3,173	27.6
Total	11,512	100.0
<i>Excluded Unknown: 37</i>	<i>0.3</i>	

% Poverty Level	Number	%
0 - 50	2,992	32.1
51 - 100	3,155	33.8
101 - 130	1,566	16.8
131 - 150	759	8.1
151 - 185	646	6.9
186 - 200	68	0.7
> 200	140	1.5
Adjunctive Eligibility	0	0.0
Total	9,326	100.0
<i>Excluded Unknown: 6</i>	<i>0.1</i>	

Program Participation ⁽³⁾	Number	%
WIC	9,332	100.0
SNAP	745	23.4
Medicaid	766	74.7
TANF	766	9.9
<i>Excluded Unknown:</i>		
<i>WIC:</i>	0	0.0
<i>SNAP:</i>	8,587	92.0
<i>Medicaid:</i>	8,566	91.8
<i>TANF:</i>	8,566	91.8

Migrant Status	Number	%
Yes	10	0.1
No	11,515	99.9
Total	11,525	100.0
<i>Excluded Unknown: 24</i>	<i>0.2</i>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Health Indicators

Summary of Health Indicators (2)				
	Nebraska Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	11,362	9.5	1,175,574	10.9
Normal Weight	11,362	43.1	1,175,574	44.6
Overweight	11,362	15.4	1,175,574	15.1
Obese	11,362	31.9	1,175,574	29.4
Maternal Weight Gain (4)				
Less Than Ideal	10,619	24.6	922,658	25.0
Ideal	10,619	34.2	922,658	32.2
Greater Than Ideal	10,619	41.2	922,658	42.8
Anemia (Low Hb/Hct) (5)				
1st Trimester	2,498	6.4	377,717	7.1
2nd Trimester	3,731	11.4	343,815	11.5
3rd Trimester	1,723	28.1	180,176	33.5
Postpartum	4,931	26.5	414,735	29.8
Parity (6)				
0	11,511	31.4	966,532	45.6
1	11,511	27.0	966,532	25.9
2	11,511	18.8	966,532	15.3
≥ 3	11,511	22.8	966,532	13.2
Interpregnancy Interval				
< 6 Months	4,800	1.7	606,104	13.0
6 - <18 Months	4,800	14.8	606,104	26.8
≥18 Months	4,800	83.5	606,104	60.3
Gestational Diabetes	0	*	230,547	3.9
Hypertension During Pregnancy	15	*	280,574	4.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI < 19.8), normal weight (BMI 19.9 to 26.0), overweight (BMI >26.0 to 29.0), obese (BMI >29.0).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Health Indicators

Summary of Health Indicators (2)				
	Nebraska Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	10,769	87.0	612,462	79.4
No Care	10,769	2.7	612,462	4.7
WIC Enrollment				
1st Trimester	11,303	26.5	1,034,046	31.3
2nd Trimester	11,303	36.6	1,034,046	35.1
3rd Trimester	11,303	17.7	1,034,046	18.2
Postpartum (3)	11,303	19.2	1,034,046	15.4
Prepregnancy Multivitamin Consumption (4)	0	*	98,170	15.7
Multivitamin Consumption During Pregnancy (5)	4	*	238,509	75.9
Smoking				
3 Months Prior to Pregnancy	11,513	34.4	895,227	26.6
During Pregnancy	8,514	19.1	826,305	15.8
Last 3 Months of Pregnancy	10,762	17.4	716,629	15.8
Postpartum	10,777	20.3	688,071	18.2
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	2,969	45.6	208,650	42.0
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	2,665	39.1	161,864	35.1
Smoking in Household				
During Pregnancy	184	100.0	572,120	20.9
Postpartum	3,270	100.0	486,636	22.6
Drinking				
3 Months Prior to Pregnancy	11,478	18.9	998,472	7.1
Last 3 Months of Pregnancy	10,748	0.2	758,070	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Health Indicators

Summary of Health Indicators (2)				
	Nebraska Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	10,440	1.1	1,039,036	1.1
Low	10,440	6.4	1,039,036	7.1
Normal	10,440	85.7	1,039,036	85.1
High	10,440	6.8	1,039,036	6.7
Preterm (4)	10,366	10.5	947,705	11.9
Full term Low Birth Weight (5)	9,255	3.2	856,467	3.7
Multiple Births	10,667	2.0	1,026,255	1.5
Breastfeeding				
Ever Breastfed	10,465	70.3	936,194	64.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.



Table 3C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Comparison of Racial and Ethnic Distribution by Contributor to Nation

Comparison of Racial and Ethnic Distribution ⁽²⁾								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Nebraska	11,549	56.9	10.7	27.9	1.5	1.8	1.1	0.0
Nation (Prior Year)	1,252,728	41.7	23.3	27.9	1.1	3.2	1.0	1.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Comparison of Age Distribution by Contributor to Nation

Comparison of Age Distribution ⁽²⁾								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Nebraska	11,549	0.3	5.8	12.3	62.4	17.9	1.3	0.1
Nation (Prior Year)	1,252,728	0.4	6.5	12.7	59.9	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Comparison of Maternal Health Indicators by Contributor to Nation

Comparison of Maternal Health Indicators ⁽²⁾														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
		Number	%		%	Number	%	%	Number	%			Number	%
Nebraska	11,362	9.5	47.3	10,619	24.6	41.2	1,723	28.1	4,931	26.5	0	*	15	*
Nation (Prior Year)	1,175,574	10.9	44.5	922,658	25.0	42.8	180,176	33.5	414,735	29.8	230,547	3.9	280,574	4.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Comparison of Maternal Behavioral Indicators by Contributor to Nation

Comparison of Maternal Behavioral Indicators ⁽²⁾									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Nebraska	10,769	87.0	2.7	11,303	26.5	0	*	4	*
Nation (Prior Year)	612,462	79.4	4.7	1,034,046	31.3	98,170	15.7	238,509	75.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Comparison of Smoking Indicators by Contributor to Nation

Comparison of Smoking Indicators (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	%	Number	%	Number	%	Number	%		
Nebraska	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
Nation (Prior Year)	895,227	26.6	716,629	15.8	161,864	35.1	572,120	20.9	486,636	22.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Comparison of Infant Health Indicators by Contributor to Nation

Comparison of Infant Health Indicators ⁽²⁾									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
Nebraska	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
Nation (Prior Year)	1,039,036	8.2	6.7	947,705	11.9	856,467	3.7	936,194	64.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 9C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska

Maternal Health Indicators by Race/Ethnicity, Age, or Education

Maternal Health Indicators by Race/Ethnicity, Age, or Education (2)

	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes		Hypertension During Pregnancy		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum		Number	%	Number	%	
							Number	%	Number	%					
Race/Ethnicity															
White, Not Hispanic	6,462	10.2	47.8	6,037	22.5	45.4	798	25.3	2,581	20.6	0	*	13	*	
Black, Not Hispanic	1,213	11.0	47.5	1,100	26.1	39.1	150	42.7	584	47.6	0	*	1	*	
Hispanic	3,183	6.5	48.2	3,021	27.8	34.3	686	26.5	1,549	28.5	0	*	1	*	
American Indian/Alaska Native	178	6.7	54.5	158	21.5	44.3	35	*	67	*	0	*	0	*	
Asian/Pacific Islander	205	27.8	17.6	194	35.6	29.4	41	*	95	*	0	*	0	*	
Multiple Races	119	12.6	38.7	108	23.1	40.7	13	*	53	*	0	*	0	*	
All Other/Unknown	2	*	*	1	*	*	0	*	2	*	0	*	0	*	
Total	11,362	9.5	47.3	10,619	24.6	41.2	1,723	28.1	4,931	26.5	0	*	15	*	
Age															
< 15 Years	33	*	*	29	*	*	2	*	14	*	0	*	0	*	
15 - 17 Years	656	17.8	25.6	601	22.1	46.3	111	32.4	293	33.1	0	*	1	*	
18 - 19 Years	1,399	13.9	32.0	1,277	20.0	47.7	213	33.3	553	35.4	0	*	2	*	
20 - 29 Years	7,098	9.4	48.2	6,650	24.4	41.1	1,035	26.5	3,067	25.7	0	*	9	*	
30 - 39 Years	2,021	4.7	61.0	1,917	28.5	36.4	338	28.7	936	21.6	0	*	3	*	
>= 40 Years	148	4.1	68.9	139	36.7	30.2	22	*	68	*	0	*	0	*	
Unknown	7	*	*	6	*	*	2	*	0	*	0	*	0	*	
Total	11,362	9.5	47.3	10,619	24.6	41.2	1,723	28.1	4,931	26.5	0	*	15	*	
Education															
< High School	3,709	9.5	43.9	3,473	28.0	37.7	695	29.8	1,695	30.7	0	*	4	*	
High School	4,518	10.2	47.6	4,211	22.8	42.3	659	29.0	1,874	25.3	0	*	7	*	
> High School	3,133	8.8	51.0	2,933	23.2	43.9	368	23.6	1,361	22.9	0	*	4	*	
Unknown	2	*	*	2	*	*	1	*	1	*	0	*	0	*	
Total	11,362	9.5	47.3	10,619	24.6	41.2	1,723	28.1	4,931	26.5	0	*	15	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 10C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education (2)									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	6,124	89.0	2.7	6,435	29.6	0	*	4	*
Black, Not Hispanic	1,115	87.4	1.1	1,219	24.5	0	*	0	*
Hispanic	3,057	83.4	3.2	3,140	21.5	0	*	0	*
American Indian/Alaska Native	160	80.6	1.3	178	20.2	0	*	0	*
Asian/Pacific Islander	200	89.0	0.5	211	19.9	0	*	0	*
Multiple Races	111	83.8	4.5	118	28.8	0	*	0	*
All Other/Unknown	2	*	*	2	*	0	*	0	*
Total	10,769	87.0	2.7	11,303	26.5	0	*	4	*
Age									
< 15 Years	30	*	*	34	*	0	*	0	*
15 - 17 Years	607	78.6	2.6	661	28.1	0	*	0	*
18 - 19 Years	1,293	84.5	3.1	1,386	29.6	0	*	1	*
20 - 29 Years	6,743	87.8	2.6	7,064	26.4	0	*	1	*
30 - 39 Years	1,951	88.8	2.9	2,006	24.1	0	*	2	*
>= 40 Years	139	86.3	1.4	145	26.9	0	*	0	*
Unknown	6	*	*	7	*	0	*	0	*
Total	10,769	87.0	2.7	11,303	26.5	0	*	4	*
Education									
< High School	3,522	82.3	3.2	3,674	24.8	0	*	1	*
High School	4,271	87.9	2.8	4,491	27.9	0	*	2	*
> High School	2,974	91.3	1.8	3,105	26.2	0	*	1	*
Unknown	2	*	*	33	*	0	*	0	*
Total	10,769	87.0	2.7	11,303	26.5	0	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 11C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Smoking Indicators by Race/Ethnicity, Age, or Education

Smoking Indicators by Race/Ethnicity, Age, or Education ⁽²⁾										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	6,550	48.9	6,121	26.2	2,171	36.2	137	100.0	2,536	100.0
Black, Not Hispanic	1,232	25.7	1,116	13.1	193	36.8	21	*	254	100.0
Hispanic	3,217	8.2	3,054	2.0	188	69.1	15	*	331	100.0
American Indian/Alaska Native	179	52.0	158	17.7	57	*	7	*	63	*
Asian/Pacific Islander	211	10.4	200	4.5	15	*	2	*	35	*
Multiple Races	122	47.5	111	19.8	40	*	2	*	50	*
All Other/Unknown	2	*	2	*	1	*	0	*	1	*
Total	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
Age										
< 15 Years	35	*	30	*	3	*	2	*	4	*
15 - 17 Years	664	26.7	606	8.9	128	59.4	17	*	217	100.0
18 - 19 Years	1,416	40.3	1,293	17.6	409	46.0	31	*	464	100.0
20 - 29 Years	7,182	37.1	6,736	19.0	1,783	37.7	102	100.0	2,144	100.0
30 - 39 Years	2,060	24.6	1,952	14.7	321	29.9	27	*	408	100.0
>= 40 Years	149	22.8	139	15.8	21	*	5	*	31	*
Unknown	7	*	6	*	0	*	0	*	2	*
Total	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
Education										
< High School	3,757	27.7	3,517	15.3	735	35.9	64	*	965	100.0
High School	4,581	41.3	4,270	21.5	1,299	37.6	86	*	1,546	100.0
> High School	3,173	32.4	2,973	14.1	631	45.6	33	*	758	100.0
Unknown	2	*	2	*	0	*	1	*	1	*
Total	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 12C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Infant Health Indicators by Race/Ethnicity, Age, or Education

Infant Health Indicators by Race/Ethnicity, Age, or Education (2)									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)						
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	5,942	7.0	7.3	5,907	10.2	5,293	2.8	5,957	66.7
Black, Not Hispanic	1,085	12.4	4.7	1,079	12.8	939	5.6	1,088	59.6
Hispanic	2,951	6.9	6.6	2,922	10.3	2,614	3.0	2,957	81.7
American Indian/Alaska Native	155	6.5	12.3	156	10.9	138	3.6	156	69.9
Asian/Pacific Islander	196	7.7	2.0	194	10.3	174	2.9	196	71.9
Multiple Races	109	5.5	5.5	106	10.4	95	*	109	58.7
All Other/Unknown	2	*	*	2	*	2	*	2	*
Total	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
Age									
< 15 Years	28	*	*	27	*	25	*	28	*
15 - 17 Years	597	6.0	4.9	594	10.8	530	3.4	597	68.0
18 - 19 Years	1,267	6.8	5.0	1,254	9.4	1,135	3.4	1,268	68.6
20 - 29 Years	6,559	6.0	6.6	6,515	9.8	5,866	2.9	6,575	70.1
30 - 39 Years	1,857	7.5	9.3	1,845	13.7	1,587	3.7	1,865	72.2
>= 40 Years	126	9.5	10.3	125	13.6	108	6.5	126	77.8
Unknown	6	*	*	6	*	4	*	6	*
Total	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
Education									
< High School	3,402	6.3	6.3	3,372	10.7	3,004	3.5	3,411	69.4
High School	4,151	6.4	5.9	4,125	10.3	3,697	3.0	4,160	67.0
> High School	2,883	6.6	8.7	2,867	10.7	2,553	3.0	2,892	75.9
Unknown	4	*	*	2	*	1	*	2	*
Total	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 13C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska

Maternal Weight Gain and Birth Outcomes by Select Health Indicators

Maternal Weight Gain and Birth Outcomes by Select Health Indicators (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal	> Ideal		Birthweight			Preterm (6)	Full Term LBW (7)		
		Low (4)	High (5)	Number	%	%				
Number	%	%	Number	%	%	Number	%	Number	%	
Prepregnancy BMI										
Underweight	1,002	29.9	25.9	985	9.8	2.2	980	12.1	861	4.3
Normal Weight	4,561	26.0	37.7	4,448	7.3	5.6	4,428	9.9	3,981	3.1
Overweight	1,633	14.7	54.1	1,584	6.9	7.2	1,574	9.3	1,422	3.1
Obese	3,399	25.7	44.4	3,301	7.3	9.6	3,273	11.5	2,892	2.9
Weight Gain										
< Ideal	-	-	-	2,419	11.8	3.3	2,403	13.7	2,067	4.9
Ideal	-	-	-	3,569	7.2	5.7	3,547	10.7	3,162	3.0
> Ideal	-	-	-	4,324	5.5	9.6	4,299	8.7	3,920	2.4
Smoking During Pregnancy										
Smokers	1,446	24.1	43.9	1,438	9.9	3.9	1,433	10.6	1,278	4.9
Non-Smokers	6,206	24.4	40.7	6,123	6.6	7.4	6,081	10.6	5,427	2.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Racial and Ethnic Distribution

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2008	11,549	56.9	10.7	27.9	1.5	1.8	1.1	0.0
2007	11,858	57.0	10.8	27.9	1.8	1.7	0.9	0.0
2006	12,457	57.2	10.8	27.8	1.8	1.7	0.6	0.0
2005	11,821	57.5	9.9	28.1	1.9	2.1	0.6	0.1
2004	11,708	58.3	9.7	27.2	1.9	2.1	0.2	0.5
2003	11,814	59.5	9.0	26.8	1.6	2.0	0.0	1.1
2002	11,401	60.9	9.4	25.4	1.8	1.7	0.0	0.9
2001	11,016	62.9	9.7	23.2	1.5	1.9	0.0	0.8
2000	10,381	64.6	10.0	21.2	1.4	2.1	0.0	0.7
1999	9,737	66.0	10.4	19.4	1.3	1.9	0.0	1.0
1998	8,945	68.8	9.0	18.3	1.2	1.9	0.0	0.8
1997	4,132	73.5	5.5	17.8	1.0	1.5	0.0	0.7
1996	7,004	71.6	9.9	14.9	1.1	1.8	0.0	0.7
1995	0	*	*	*	*	*	*	*
1994	7,538	74.6	11.3	11.1	1.2	1.6	0.0	0.2
1993	5,955	76.9	10.7	9.4	1.3	1.6	0.0	0.1
1992	6,190	75.9	12.1	8.6	1.8	1.5	0.0	0.1
1991	6,208	76.1	12.8	7.4	2.2	1.4	0.0	0.1
1990	2,259	79.6	11.2	6.0	1.6	1.4	0.0	0.0
1989	2,452	78.1	12.7	5.5	1.9	1.3	0.0	0.6
1988	3,298	76.1	14.6	4.9	2.5	1.6	0.0	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Racial and Ethnic Distribution

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1987	3,790	75.5	16.2	4.7	2.1	1.3	0.0	0.2
1986	2,421	77.9	14.5	4.2	1.7	1.1	0.0	0.6
1985	724	78.9	10.9	2.6	1.5	5.7	0.0	0.4
1984	2,437	71.3	18.8	4.6	2.2	2.1	0.0	1.1
1983	2,492	71.8	18.9	4.8	2.0	1.6	0.0	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2008	11,549	0.3	5.8	12.3	62.4	17.9	1.3	0.1
2007	11,858	0.3	6.0	12.2	63.4	17.1	1.0	0.0
2006	12,457	0.3	5.5	12.3	62.7	17.9	1.2	0.1
2005	11,821	0.3	6.2	12.0	62.7	17.7	1.1	0.1
2004	11,708	0.3	6.2	12.9	62.0	17.3	1.2	0.1
2003	11,814	0.3	6.3	13.3	62.4	16.6	1.0	0.0
2002	11,401	0.3	6.7	14.5	60.7	16.9	1.0	0.0
2001	11,016	0.3	7.2	13.8	61.3	16.3	1.0	0.0
2000	10,381	0.4	7.3	15.0	60.0	16.1	1.2	0.0
1999	9,737	0.4	8.3	14.6	60.2	15.6	0.9	0.0
1998	8,945	0.5	8.3	15.0	59.9	15.4	0.9	0.0
1997	4,132	0.6	9.3	14.8	57.6	16.6	1.1	0.0
1996	7,004	0.6	8.6	14.7	60.0	15.3	1.0	0.0
1995	0	*	*	*	*	*	*	*
1994	7,538	0.6	8.3	14.6	59.4	16.5	0.6	0.0
1993	5,955	0.6	7.6	14.0	61.4	15.8	0.6	0.0
1992	6,190	0.5	7.7	14.0	62.5	14.5	0.7	0.0
1991	6,208	0.5	7.3	14.4	63.1	14.0	0.6	0.0
1990	2,259	0.4	6.8	14.5	62.3	15.8	0.2	0.0
1989	2,452	0.7	9.1	14.0	61.7	13.9	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1988	3,298	0.7	8.5	11.9	64.9	13.5	0.5	0.1
1987	3,790	0.6	8.1	14.6	64.5	11.7	0.4	0.1
1986	2,421	0.6	6.9	13.8	66.2	11.9	0.5	0.2
1985	724	0.6	7.7	12.6	64.9	13.5	0.7	0.0
1984	2,437	0.7	9.0	18.1	60.9	9.7	0.3	1.3
1983	2,492	0.6	9.9	18.5	60.9	9.0	0.2	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2008	11,362	9.5	47.3	10,619	24.6	41.2	1,723	28.1	4,931	26.5
2007	11,646	9.7	45.9	10,923	24.4	40.8	1,631	33.7	5,319	28.6
2006	12,172	10.3	44.5	11,238	24.5	40.6	1,321	28.3	5,413	27.8
2005	11,582	10.1	44.3	10,627	24.9	41.3	1,624	27.8	5,655	29.3
2004	11,510	10.9	43.8	10,480	24.1	42.5	1,509	26.8	5,704	28.4
2003	11,553	11.4	42.0	10,470	28.7	39.7	1,567	25.6	5,606	25.9
2002	11,120	11.7	41.7	10,047	29.9	38.9	1,671	23.9	5,371	24.9
2001	10,810	12.1	40.7	9,886	29.8	38.0	1,607	26.8	4,723	23.4
2000	10,175	13.6	39.5	9,344	29.4	37.8	1,640	26.5	4,499	26.4
1999	9,524	13.3	39.5	8,658	28.0	39.0	1,831	26.1	4,174	23.3
1998	8,687	13.9	38.6	8,010	27.7	38.4	1,595	24.7	3,680	24.0
1997	3,881	14.0	37.0	3,431	25.4	37.9	697	22.2	1,323	19.3
1996	6,742	15.4	35.8	6,471	25.5	38.9	1,183	26.3	2,776	21.1
1995	0	*	*	0	*	*	0	*	0	*
1994	7,137	16.4	33.8	6,356	26.6	37.5	1,330	25.0	3,151	19.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1993	5,716	16.3	33.5	5,318	25.7	38.2	1,065	22.3	2,871	16.1
1992	5,777	17.0	32.3	5,335	28.1	36.8	1,087	22.3	3,165	17.5
1991	5,625	17.4	31.3	5,247	28.7	35.9	1,096	21.8	3,546	16.1
1990	1,901	19.4	28.5	1,787	25.8	29.4	394	19.5	1,416	12.9
1989	2,425	17.6	30.8	0	*	*	622	21.4	588	14.1
1988	3,278	17.4	31.5	0	*	*	982	19.3	258	16.7
1987	3,756	17.6	29.5	0	*	*	1,079	16.6	1,491	16.0
1986	2,368	19.6	29.9	0	*	*	665	15.8	433	8.1
1985	602	20.9	25.9	0	*	*	205	12.7	10	*
1984	2,391	20.8	27.2	0	*	*	705	19.4	1,637	9.8
1983	2,437	22.7	25.0	0	*	*	770	23.9	1,614	12.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2008	10,769	87.0	2.7	11,303	26.5
2007	10,998	88.4	1.9	11,575	27.0
2006	11,298	88.3	1.9	12,196	30.0
2005	10,692	88.5	1.7	11,574	26.7
2004	10,536	88.4	1.7	10,992	29.2
2003	10,574	87.5	2.3	11,587	27.9
2002	10,125	87.4	1.8	11,082	27.4
2001	9,968	87.3	1.7	10,826	25.7
2000	9,412	86.1	1.9	10,177	25.5
1999	8,745	85.7	2.3	8,168	30.0
1998	8,039	84.9	2.1	7,784	28.6
1997	3,205	82.1	1.6	3,919	27.1
1996	5,652	82.4	1.3	6,724	25.9
1995	0	*	*	0	*
1994	4,353	77.3	2.2	6,056	26.4
1993	2,951	76.8	2.7	4,762	26.0
1992	90	*	*	4,858	20.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
1991	0	*	*	4,646	18.6
1990	0	*	*	1,613	16.4
1989	0	*	*	2,242	21.9
1988	0	*	*	3,140	19.5
1987	0	*	*	3,633	14.4
1986	0	*	*	2,295	12.9
1985	0	*	*	652	18.3
1984	0	*	*	2,320	18.0
1983	0	*	*	2,304	16.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Smoking Indicators

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2008	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
2007	11,750	33.1	10,996	16.3	2,472	41.4	255	100.0	3,320	100.0
2006	12,259	33.9	11,279	16.8	2,640	40.2	261	100.0	3,502	100.0
2005	11,674	33.2	6,175	16.1	1,131	42.3	347	99.7	3,372	100.0
2004	11,564	33.8	2,726	18.0	242	40.5	421	81.9	5,299	66.4
2003	11,665	33.8	1,920	20.0	15	*	1,085	36.4	10,641	32.3
2002	11,224	34.2	2,133	19.4	33	*	1,088	41.2	10,212	32.7
2001	10,913	35.4	2,189	19.8	27	*	920	33.8	10,027	33.4
2000	10,281	35.6	2,257	21.4	13	*	725	40.0	9,472	34.5
1999	9,668	36.4	2,085	21.5	50	*	655	35.0	8,817	34.1
1998	8,867	37.3	2,115	20.9	188	45.2	339	36.0	8,176	34.7
1997	4,015	37.6	1,095	21.0	128	25.8	205	4.4	1,435	35.1
1996	6,737	38.7	2,027	26.7	138	18.8	1	*	0	*
1995	0	*	0	*	0	*	0	*	0	*
1994	7,041	38.8	1,869	29.2	144	11.1	0	*	0	*
1993	5,535	40.6	1,578	26.5	193	19.7	0	*	0	*
1992	5,592	40.6	1,473	29.1	121	18.2	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Smoking Indicators

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%		
1991	5,154	43.1	1,566	31.7	127	16.5	0	*	0	*
1990	1,044	42.7	570	36.5	26	*	0	*	0	*
1989	0	*	1	*	0	*	0	*	0	*
1988	0	*	0	*	0	*	0	*	0	*
1987	0	*	0	*	0	*	0	*	0	*
1986	0	*	0	*	0	*	0	*	0	*
1985	0	*	0	*	0	*	0	*	0	*
1984	0	*	0	*	0	*	0	*	0	*
1983	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2008	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
2007	10,795	7.3	7.3	10,611	10.8	9,440	3.1	10,807	69.7
2006	11,118	7.1	7.0	10,937	11.6	9,656	3.0	11,105	69.4
2005	10,520	7.1	7.7	10,377	10.7	9,249	3.1	8,911	99.9
2004	10,370	7.0	7.6	10,249	10.2	9,184	3.2	9,146	64.6
2003	10,397	7.1	7.8	10,238	11.0	9,094	3.2	9,179	62.9
2002	9,996	7.3	8.5	9,765	11.1	8,663	2.9	8,901	60.2
2001	9,778	6.9	8.1	9,653	10.9	8,583	3.1	8,928	58.5
2000	9,202	7.2	8.2	9,056	10.6	8,082	3.2	8,584	58.3
1999	8,499	6.9	8.6	8,299	10.8	7,347	2.9	7,914	53.9
1998	7,823	6.5	8.9	7,378	11.0	6,535	3.0	7,283	50.1
1997	3,491	7.9	8.9	3,300	10.9	2,933	3.2	2,368	61.8
1996	6,598	6.5	9.4	6,363	9.9	5,722	2.7	6,345	48.3
1995	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1994	6,615	6.9	10.0	5,221	8.8	4,754	3.1	6,237	50.2
1993	5,478	6.5	9.8	4,346	9.7	3,921	2.6	5,184	49.8
1992	5,631	6.2	10.7	4,388	8.9	3,984	2.8	5,079	47.0
1991	5,731	6.5	9.8	4,287	8.3	3,923	3.2	4,714	49.2
1990	2,157	6.8	9.9	1,555	8.4	1,416	3.0	1,364	62.8
1989	2,445	7.9	9.9	2,208	11.7	1,950	4.0	711	100.0
1988	3,286	6.0	10.1	3,073	10.3	2,755	3.5	605	100.0
1987	3,768	5.8	10.9	3,575	12.0	3,138	3.2	712	100.0
1986	2,396	6.5	11.4	2,248	11.0	1,996	3.2	675	100.0
1985	719	6.0	10.6	638	10.2	573	3.1	247	100.0
1984	2,400	6.2	10.4	2,287	11.4	2,021	3.3	754	100.0
1983	2,462	6.4	9.6	2,259	11.4	2,000	3.9	794	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2008	6,462	10.2	47.8	6,037	22.5	45.4	798	25.3	2,581	20.6
	2007	6,633	10.4	46.5	6,206	21.1	45.5	830	28.8	2,767	23.0
	2006	6,974	11.1	44.5	6,426	21.4	45.8	732	22.7	2,836	19.6
	2005	6,667	10.9	44.6	6,138	22.3	45.4	872	24.3	3,057	22.0
	2004	6,721	12.3	43.6	6,136	20.4	47.3	890	21.5	3,043	20.6
	2003	6,879	13.2	41.0	6,215	23.6	45.1	919	21.4	3,070	19.8
	2002	6,763	13.7	41.5	6,099	24.6	44.5	973	20.6	3,093	19.4
	2001	6,796	13.8	39.8	6,255	26.1	42.0	955	21.3	2,808	18.9
	2000	6,595	15.0	38.7	6,073	26.3	41.5	1,023	22.6	2,775	21.7
	1999	6,305	14.4	39.3	5,774	26.0	41.3	1,134	23.5	2,713	18.5
Black, Not Hispanic	2008	1,213	11.0	47.5	1,100	26.1	39.1	150	42.7	584	47.6
	2007	1,260	11.0	46.5	1,146	25.8	37.5	169	56.8	673	45.6
	2006	1,315	12.3	44.2	1,151	26.6	37.1	156	52.6	741	44.1
	2005	1,137	10.7	45.9	972	26.4	40.3	126	37.3	704	48.7
	2004	1,113	11.1	44.9	931	23.4	41.7	120	47.5	661	48.6
	2003	1,024	11.1	45.7	879	25.6	41.4	100	49.0	671	42.2
	2002	1,035	10.3	43.9	898	43.0	30.2	176	45.5	687	38.4
	2001	1,049	10.7	45.9	905	46.0	27.0	149	49.7	530	36.2
	2000	1,014	13.6	41.7	904	45.6	27.1	164	51.8	575	45.7
	1999	985	13.5	39.8	841	38.5	31.4	211	46.9	439	37.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2008	3,183	6.5	48.2	3,021	27.8	34.3	686	26.5	1,549	28.5
	2007	3,259	6.3	46.1	3,112	29.6	33.3	561	34.6	1,663	31.2
	2006	3,383	7.0	46.5	3,200	29.9	31.8	372	28.8	1,629	33.6
	2005	3,257	7.1	45.1	3,059	29.4	34.1	545	29.9	1,657	33.9
	2004	3,131	6.4	45.9	2,926	31.7	33.7	439	32.6	1,746	33.7
	2003	3,106	6.4	44.8	2,892	40.4	29.0	474	27.4	1,568	30.2
	2002	2,845	6.0	43.1	2,614	37.7	29.4	456	22.6	1,353	30.0
	2001	2,513	7.5	42.6	2,316	32.5	32.4	429	29.6	1,179	26.5
	2000	2,140	7.2	43.6	1,988	28.6	33.3	384	25.8	950	27.5
	1999	1,837	7.5	42.6	1,686	28.1	37.1	404	23.5	838	32.0
American Indian/Alaska Native	2008	178	6.7	54.5	158	21.5	44.3	35	*	67	*
	2007	200	12.5	48.0	182	27.5	38.5	33	*	83	*
	2006	216	5.6	47.7	200	18.5	42.5	24	*	97	*
	2005	217	8.3	47.5	182	21.4	44.5	39	*	96	*
	2004	220	11.4	48.6	191	25.7	44.0	29	*	95	*
	2003	185	7.0	47.6	158	20.9	46.2	30	*	96	*
	2002	192	12.0	52.1	164	23.8	45.1	31	*	79	*
	2001	160	8.1	45.0	143	28.7	37.8	30	*	61	*
	2000	138	8.7	40.6	114	33.3	43.0	24	*	54	*
	1999	123	11.4	42.3	97	*	*	22	*	44	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2008	205	27.8	17.6	194	35.6	29.4	41	*	95	*
	2007	196	29.6	14.8	187	32.1	28.3	27	*	94	*
	2006	211	30.8	14.7	194	32.5	27.8	27	*	80	*
	2005	238	29.0	15.5	219	29.7	25.1	36	*	110	19.1
	2004	244	29.1	15.6	223	26.9	31.4	25	*	117	20.5
	2003	238	32.4	12.6	221	35.3	19.0	32	*	132	23.5
	2002	190	28.9	11.6	182	37.9	23.1	30	*	108	20.4
	2001	207	23.7	19.3	191	42.4	27.7	36	*	99	*
	2000	214	35.0	15.9	198	53.0	20.2	34	*	100	26.0
	1999	183	32.8	16.9	175	41.1	24.0	36	*	88	*
Multiple Races	2008	119	12.6	38.7	108	23.1	40.7	13	*	53	*
	2007	98	*	*	90	*	*	11	*	39	*
	2006	73	*	*	67	*	*	10	*	30	*
	2005	64	*	*	57	*	*	6	*	31	*
	2004	29	*	*	29	*	*	3	*	13	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2008	2	*	*	1	*	*	0	*	2	*
	2007	0	*	*	0	*	*	0	*	0	*
	2006	0	*	*	0	*	*	0	*	0	*
	2005	2	*	*	0	*	*	0	*	0	*
	2004	52	*	*	44	*	*	3	*	29	*
	2003	121	9.9	45.5	105	32.4	34.3	12	*	69	*
	2002	95	*	*	90	*	*	5	*	51	*
	2001	85	*	*	76	*	*	8	*	46	*
	2000	74	*	*	67	*	*	11	*	45	*
	1999	91	*	*	85	*	*	24	*	52	*
Total All Race/Ethnic Groups	2008	11,362	9.5	47.3	10,619	24.6	41.2	1,723	28.1	4,931	26.5
	2007	11,646	9.7	45.9	10,923	24.4	40.8	1,631	33.7	5,319	28.6
	2006	12,172	10.3	44.5	11,238	24.5	40.6	1,321	28.3	5,413	27.8
	2005	11,582	10.1	44.3	10,627	24.9	41.3	1,624	27.8	5,655	29.3
	2004	11,510	10.9	43.8	10,480	24.1	42.5	1,509	26.8	5,704	28.4
	2003	11,553	11.4	42.0	10,470	28.7	39.7	1,567	25.6	5,606	25.9
	2002	11,120	11.7	41.7	10,047	29.9	38.9	1,671	23.9	5,371	24.9
	2001	10,810	12.1	40.7	9,886	29.8	38.0	1,607	26.8	4,723	23.4
	2000	10,175	13.6	39.5	9,344	29.4	37.8	1,640	26.5	4,499	26.4
	1999	9,524	13.3	39.5	8,658	28.0	39.0	1,831	26.1	4,174	23.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2008	6,124	89.0	2.7	6,435	29.6
	2007	6,249	89.6	1.9	6,604	29.3
	2006	6,461	90.1	1.5	6,971	31.8
	2005	6,170	90.0	1.6	6,652	29.4
	2004	6,173	89.7	1.6	6,356	31.3
	2003	6,284	89.3	2.1	6,897	29.7
	2002	6,151	89.6	1.6	6,733	29.7
	2001	6,316	88.8	1.6	6,799	27.8
	2000	6,109	88.3	1.7	6,566	27.2
	1999	5,828	87.7	2.1	5,395	32.2
Black, Not Hispanic	2008	1,115	87.4	1.1	1,219	24.5
	2007	1,153	87.7	0.6	1,258	21.5
	2006	1,155	86.5	1.0	1,328	21.5
	2005	975	86.9	0.5	1,147	19.2
	2004	933	88.9	0.2	1,092	21.6
	2003	889	87.5	1.3	1,049	22.1
	2002	907	84.9	0.8	1,042	23.0
	2001	911	84.7	0.7	1,051	23.8
	2000	910	81.9	1.2	1,021	20.0
	1999	849	87.0	0.6	819	23.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2008	3,057	83.4	3.2	3,140	21.5
	2007	3,128	86.9	2.4	3,216	24.6
	2006	3,214	85.7	3.0	3,389	29.8
	2005	3,084	86.3	2.0	3,244	23.9
	2004	2,945	85.9	2.3	3,027	27.6
	2003	2,914	83.9	3.1	3,102	25.7
	2002	2,630	83.7	2.4	2,831	23.4
	2001	2,330	84.8	2.4	2,519	21.2
	2000	2,014	81.7	2.8	2,161	23.0
	1999	1,707	78.6	3.9	1,612	25.7
American Indian/Alaska Native	2008	160	80.6	1.3	178	20.2
	2007	188	76.1	1.6	204	25.5
	2006	204	80.9	1.5	223	22.0
	2005	185	81.6	1.6	219	22.4
	2004	189	76.7	5.3	214	25.2
	2003	159	79.9	4.4	181	24.9
	2002	162	79.6	2.5	191	23.0
	2001	143	82.5	2.8	162	22.2
	2000	116	75.0	3.4	141	20.6
	1999	98	*	*	104	28.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2008	200	89.0	0.5	211	19.9
	2007	189	92.6	1.1	195	23.6
	2006	195	89.7	1.5	212	32.5
	2005	220	89.5	3.2	243	27.2
	2004	223	91.0	0.9	228	29.8
	2003	222	89.6	0.5	235	23.8
	2002	183	89.1	1.1	192	20.8
	2001	191	84.3	1.6	211	22.7
	2000	197	87.3	1.5	215	25.1
1999	176	89.8	1.7	159	26.4	
Multiple Races	2008	111	83.8	4.5	118	28.8
	2007	91	*	*	98	*
	2006	69	*	*	73	*
	2005	58	*	*	66	*
	2004	29	*	*	28	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
	2000	0	*	*	0	*
1999	0	*	*	0	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2008	2	*	*	2	*
	2007	0	*	*	0	*
	2006	0	*	*	0	*
	2005	0	*	*	3	*
	2004	44	*	*	47	*
	2003	106	86.8	0.9	123	39.0
	2002	92	*	*	93	*
	2001	77	*	*	84	*
	2000	66	*	*	73	*
	1999	87	*	*	79	*
Total All Race/Ethnic Groups	2008	10,769	87.0	2.7	11,303	26.5
	2007	10,998	88.4	1.9	11,575	27.0
	2006	11,298	88.3	1.9	12,196	30.0
	2005	10,692	88.5	1.7	11,574	26.7
	2004	10,536	88.4	1.7	10,992	29.2
	2003	10,574	87.5	2.3	11,587	27.9
	2002	10,125	87.4	1.8	11,082	27.4
	2001	9,968	87.3	1.7	10,826	25.7
	2000	9,412	86.1	1.9	10,177	25.5
	1999	8,745	85.7	2.3	8,168	30.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2008 Pregnancy Nutrition Surveillance (1)

Nebraska
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2008	6,550	48.9	6,121	26.2	2,171	36.2	137	100.0	2,536	100.0
	2007	6,692	47.1	6,247	24.9	2,026	38.0	187	100.0	2,551	100.0
	2006	7,020	48.4	6,450	25.7	2,181	36.9	181	100.0	2,722	100.0
	2005	6,722	46.7	3,599	23.8	933	38.5	235	100.0	2,608	100.0
	2004	6,749	47.8	1,707	25.0	194	36.1	293	86.3	3,587	74.6
	2003	6,945	48.2	1,313	26.7	14	*	659	44.3	6,319	42.3
	2002	6,825	48.1	1,395	25.6	28	*	666	48.9	6,199	41.5
	2001	6,860	48.1	1,499	24.9	21	*	533	40.7	6,350	41.9
	2000	6,646	47.6	1,548	28.2	12	*	441	47.2	6,141	43.0
	1999	6,391	47.5	1,512	25.7	39	*	372	43.8	5,866	40.9
Black, Not Hispanic	2008	1,232	25.7	1,116	13.1	193	36.8	21	*	254	100.0
	2007	1,274	22.4	1,156	10.3	157	48.4	31	*	261	100.0
	2006	1,324	22.5	1,155	10.0	155	47.7	35	*	238	100.0
	2005	1,146	26.4	615	12.2	72	*	53	*	229	100.0
	2004	1,120	23.1	199	14.1	17	*	55	*	437	61.8
	2003	1,040	22.8	133	12.0	0	*	154	32.5	892	27.1
	2002	1,051	21.2	194	14.9	2	*	145	35.9	912	29.2
	2001	1,061	23.5	200	17.0	3	*	149	29.5	912	27.4
	2000	1,028	21.8	216	10.6	1	*	102	41.2	917	24.2
	1999	1,001	19.3	172	19.2	1	*	125	21.6	859	25.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2008 Pregnancy Nutrition Surveillance (1)
Nebraska

Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2008	3,217	8.2	3,054	2.0	188	69.1	15	*	331	100.0
	2007	3,281	8.9	3,125	2.3	201	62.2	23	*	351	100.0
	2006	3,412	8.5	3,208	2.1	191	67.5	32	*	381	100.0
	2005	3,278	8.7	1,689	2.1	89	*	35	*	386	100.0
	2004	3,149	8.3	686	3.4	22	*	49	*	1,015	39.8
	2003	3,135	6.7	383	2.6	0	*	213	13.6	2,941	12.5
	2002	2,870	7.4	458	2.8	1	*	235	21.7	2,654	13.3
	2001	2,536	7.8	407	3.9	2	*	193	15.5	2,352	13.1
	2000	2,178	8.6	392	3.3	0	*	136	18.4	2,032	14.1
	1999	1,872	9.3	312	4.2	6	*	124	18.5	1,728	16.3
American Indian/Alaska Native	2008	179	52.0	158	17.7	57	*	7	*	63	*
	2007	206	45.1	188	14.4	53	*	8	*	80	*
	2006	219	49.3	203	16.7	72	*	7	*	90	*
	2005	220	40.9	112	17.0	25	*	14	*	73	*
	2004	220	52.3	44	*	7	*	14	*	114	70.2
	2003	185	53.5	28	*	1	*	27	*	160	48.8
	2002	190	47.9	35	*	2	*	30	*	168	42.9
	2001	162	45.7	32	*	0	*	20	*	144	35.4
	2000	140	38.6	32	*	0	*	22	*	117	40.2
	1999	124	46.8	25	*	2	*	21	*	98	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2008 Pregnancy Nutrition Surveillance (1)

Nebraska
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2008	211	10.4	200	4.5	15	*	2	*	35	*
	2007	198	10.6	189	3.7	9	*	4	*	40	*
	2006	211	14.2	195	5.1	18	*	3	*	44	*
	2005	241	11.2	136	1.5	5	*	3	*	57	*
	2004	244	10.7	64	*	2	*	7	*	101	56.4
	2003	239	8.8	51	*	0	*	17	*	222	23.9
	2002	192	8.3	42	*	0	*	8	*	186	28.5
	2001	208	14.9	35	*	0	*	17	*	192	29.7
	2000	215	10.2	53	*	0	*	17	*	198	28.8
	1999	186	15.6	44	*	1	*	8	*	178	26.4
Multiple Races	2008	122	47.5	111	19.8	40	*	2	*	50	*
	2007	99	*	91	*	26	*	2	*	37	*
	2006	73	*	68	*	23	*	3	*	27	*
	2005	65	*	24	*	7	*	6	*	19	*
	2004	29	*	11	*	0	*	0	*	15	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2008 Pregnancy Nutrition Surveillance (1)

Nebraska
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2008	2	*	2	*	1	*	0	*	1	*
	2007	0	*	0	*	0	*	0	*	0	*
	2006	0	*	0	*	0	*	0	*	0	*
	2005	2	*	0	*	0	*	1	*	0	*
	2004	53	*	15	*	0	*	3	*	30	*
	2003	121	18.2	12	*	0	*	15	*	107	24.3
	2002	96	*	9	*	0	*	4	*	93	*
	2001	86	*	16	*	1	*	8	*	77	*
	2000	74	*	16	*	0	*	7	*	67	*
	1999	94	*	20	*	1	*	5	*	88	*
Total All Race/Ethnic Groups	2008	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
	2007	11,750	33.1	10,996	16.3	2,472	41.4	255	100.0	3,320	100.0
	2006	12,259	33.9	11,279	16.8	2,640	40.2	261	100.0	3,502	100.0
	2005	11,674	33.2	6,175	16.1	1,131	42.3	347	99.7	3,372	100.0
	2004	11,564	33.8	2,726	18.0	242	40.5	421	81.9	5,299	66.4
	2003	11,665	33.8	1,920	20.0	15	*	1,085	36.4	10,641	32.3
	2002	11,224	34.2	2,133	19.4	33	*	1,088	41.2	10,212	32.7
	2001	10,913	35.4	2,189	19.8	27	*	920	33.8	10,027	33.4
	2000	10,281	35.6	2,257	21.4	13	*	725	40.0	9,472	34.5
	1999	9,668	36.4	2,085	21.5	50	*	655	35.0	8,817	34.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
White, Not Hispanic	2008	5,942	7.0	7.3	5,907	10.2	5,293	2.8	5,957	66.7
	2007	6,125	7.0	7.7	6,037	10.7	5,382	2.8	6,133	66.2
	2006	6,358	6.8	7.4	6,252	11.3	5,537	2.7	6,353	65.8
	2005	6,076	6.9	8.2	5,993	10.5	5,352	2.8	5,129	99.9
	2004	6,074	6.8	8.2	5,989	10.6	5,353	2.7	5,330	59.9
	2003	6,169	7.0	8.4	6,079	11.4	5,376	3.0	5,456	57.7
	2002	6,051	7.3	8.7	5,918	11.5	5,219	2.8	5,404	55.5
	2001	6,190	6.8	8.4	6,099	10.9	5,425	3.0	5,649	54.8
	2000	5,963	6.9	8.7	5,854	10.2	5,247	2.7	5,571	57.2
	1999	5,656	6.4	9.0	5,523	10.3	4,916	2.5	5,235	52.0
Black, Not Hispanic	2008	1,085	12.4	4.7	1,079	12.8	939	5.6	1,088	59.6
	2007	1,140	12.3	3.8	1,123	12.6	977	5.6	1,146	60.1
	2006	1,156	11.4	4.2	1,142	14.2	980	5.4	1,151	61.5
	2005	964	11.1	4.9	952	13.2	825	5.3	804	100.0
	2004	929	10.3	5.3	932	10.3	829	5.2	890	53.8
	2003	881	9.5	4.0	873	10.9	775	4.1	842	56.5
	2002	901	11.1	5.3	881	12.9	767	4.2	871	49.7
	2001	904	10.3	4.3	896	11.7	790	4.9	862	43.2
	2000	903	12.2	3.2	894	13.6	770	6.8	868	43.3
	1999	837	12.7	4.7	819	14.2	695	6.3	809	35.6

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
Hispanic	2008	2,951	6.9	6.6	2,922	10.3	2,614	3.0	2,957	81.7
	2007	3,068	6.0	7.4	2,998	10.5	2,680	2.6	3,066	81.5
	2006	3,142	6.0	7.3	3,084	11.2	2,737	2.8	3,140	79.8
	2005	3,021	6.3	7.3	2,975	10.5	2,657	2.9	2,588	100.0
	2004	2,891	6.4	7.4	2,856	9.4	2,583	3.4	2,494	79.1
	2003	2,869	6.3	7.7	2,816	9.9	2,533	3.2	2,459	77.8
	2002	2,610	6.0	8.9	2,550	9.7	2,300	2.8	2,243	76.4
	2001	2,279	5.6	8.4	2,259	10.1	2,025	2.6	2,046	75.3
	2000	1,967	5.8	9.0	1,947	10.0	1,750	2.9	1,797	69.6
	1999	1,661	5.7	9.0	1,620	10.7	1,440	3.0	1,550	69.5
American Indian/Alaska Native	2008	155	6.5	12.3	156	10.9	138	3.6	156	69.9
	2007	184	6.0	14.7	179	11.7	158	1.9	183	63.9
	2006	202	6.9	8.4	201	12.9	174	4.0	201	64.7
	2005	184	4.3	15.2	183	11.5	162	0.6	148	100.0
	2004	190	8.4	12.6	186	12.9	162	2.5	171	54.4
	2003	156	5.8	12.2	152	14.5	130	1.5	138	51.4
	2002	158	3.2	15.2	149	9.4	135	0.7	146	54.1
	2001	141	7.8	12.1	140	12.9	121	3.3	131	53.4
	2000	112	8.0	13.4	111	12.6	97	*	103	56.3
	1999	91	*	*	85	*	73	*	82	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2008	196	7.7	2.0	194	10.3	174	2.9	196	71.9
	2007	188	9.0	6.9	186	10.2	166	3.6	189	58.7
	2006	192	9.4	5.2	190	12.6	166	2.4	192	70.3
	2005	217	7.4	5.1	216	7.9	199	4.5	191	100.0
	2004	218	6.4	3.7	218	10.6	194	3.6	197	65.0
	2003	213	8.5	4.7	213	12.2	186	5.4	189	48.1
	2002	184	9.8	4.3	179	8.4	163	5.5	164	47.6
	2001	189	6.3	8.5	187	15.0	159	1.3	172	52.3
	2000	192	7.3	5.7	187	12.3	164	3.0	182	48.4
	1999	169	5.9	6.5	168	8.3	153	2.6	161	52.2
Multiple Races	2008	109	5.5	5.5	106	10.4	95	*	109	58.7
	2007	90	*	*	88	*	77	*	90	*
	2006	68	*	*	68	*	62	*	68	*
	2005	58	*	*	58	*	54	*	51	*
	2004	29	*	*	28	*	27	*	27	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2008	2	*	*	2	*	2	*	2	*
	2007	0	*	*	0	*	0	*	0	*
	2006	0	*	*	0	*	0	*	0	*
	2005	0	*	*	0	*	0	*	0	*
	2004	39	*	*	40	*	36	*	37	*
	2003	109	9.2	9.2	105	9.5	94	*	95	*
	2002	92	*	*	88	*	79	*	73	*
	2001	75	*	*	72	*	63	*	68	*
	2000	65	*	*	63	*	54	*	63	*
	1999	85	*	*	84	*	70	*	77	*
Total All Race/Ethnic Groups	2008	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
	2007	10,795	7.3	7.3	10,611	10.8	9,440	3.1	10,807	69.7
	2006	11,118	7.1	7.0	10,937	11.6	9,656	3.0	11,105	69.4
	2005	10,520	7.1	7.7	10,377	10.7	9,249	3.1	8,911	99.9
	2004	10,370	7.0	7.6	10,249	10.2	9,184	3.2	9,146	64.6
	2003	10,397	7.1	7.8	10,238	11.0	9,094	3.2	9,179	62.9
	2002	9,996	7.3	8.5	9,765	11.1	8,663	2.9	8,901	60.2
	2001	9,778	6.9	8.1	9,653	10.9	8,583	3.1	8,928	58.5
	2000	9,202	7.2	8.2	9,056	10.6	8,082	3.2	8,584	58.3
	1999	8,499	6.9	8.6	8,299	10.8	7,347	2.9	7,914	53.9

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

