


INJURY REPORT, 1999-2003

**Lincoln/Lancaster Health Department Coverage Area**  
Lancaster County

**Summary**

Injury is a public health problem in Lancaster County. Overall, injury is the fourth leading cause of death (*Table D1*). From 1999 to 2003, in addition to 568 injury deaths, nearly 5,800 residents in Lancaster County were hospitalized and more than 104,000 residents were treated at hospitals for injuries (*Figure A*). Overall, suicide was the leading cause of injury death. Motor vehicle crashes were second, followed by falls. In contrast to deaths, falls were the leading cause of injury-related hospital discharges and struck by/against injuries were second, followed by motor vehicle crashes, overexertion, and cutting/piercing.

**Figure A: Injury Deaths, Hospital Inpatients and Outpatients  
Lancaster County, NE, 1999-2003**



The pattern of injury varied by age and gender. Injury had a greater impact on children and young adults. Of all deaths among those age 15-29, most were due to injuries. Therefore, more years of potential life were lost due to injuries. Males were almost twice as likely as females to die from injuries. Additionally, teens and young adults were more likely to die from motor vehicle crashes, while older adults were more likely to die from fall-related injuries.

The following table highlights Lancaster County's progress towards the 2010 Nebraska Health Goals and Objectives related to deaths due to unintentional injury, falls, motor vehicle crashes, suicide, and homicide (*Table A*).

**Table A: Injury Death Rates, Lancaster County, NE 2003**

Cause of Death	Death Rate 2003	2010 Nebraska Objective (rate per 100,000 population)
Unintentional Injury	28.7	19.4
Fall	6.9	3.5
Motor Vehicle Crash	11.2	12.0
Suicide	10.5	8.2
Homicide	1.4	2.0

## Injury Deaths

### Overall

- From 1999 to 2003, injuries, including unintentional and intentional, were the fourth leading cause of death in Lancaster County (*Table D1*).

**Table D1: Five Leading Causes of Death,  
Lancaster County, NE, 1999-2003**

Rank	Cause	Total Number of Death
1	Cancer	2,020
2	Heart Disease	1,970
3	Stroke	545
4	Injury	538
5	Chronic Lung Disease	427

Source: DHHS Death Certificate Data 1999-2003

- An average of 108 residents died from injury each year from 1999 to 2003 (*Table D2*).
- Deaths due to injury made up 6.5% of total deaths in Lancaster County, compared to 6% of deaths in Nebraska.
- More than half of the deaths among 15-29 year olds were due to injuries in Lancaster County (*Table D2*).


**Table D2: Injury Deaths as a Percent of All Deaths by Age,  
Lancaster County, NE, 1999-2003**

Age Group	Annual Average		All Deaths	Percent of Death due to Injury
	Injury Deaths	Injury Deaths		
<1	4	1	120	3.3
1-4	7	1	18	38.9
5-9	3	1	9	33.3
10-14	4	1	17	23.5
15-19	30	6	38	78.9
20-24	50	10	62	80.6
25-29	28	6	50	56.0
30-34	38	8	79	48.1
35-39	46	9	116	39.7
40-44	39	8	164	23.8
45-49	52	10	235	22.1
50-54	30	6	276	10.9
55-59	19	4	286	6.6
60-64	12	2	403	3.0
65-74	29	6	1,319	2.2
75-84	55	11	2,378	2.3
85+	92	18	2,757	3.3
<b>Total</b>	<b>538</b>	<b>108</b>	<b>8,327</b>	<b>6.5</b>

Source: DHHS Death Certificate Data 1999-2003

- Years of Potential Life Lost (YPLL) from unintentional injury ranked second (13.6%), following that from cancer (22.8%). Suicide ranked 4<sup>th</sup> at 7.0% (*Figure D1*).

**Figure D1: Years of Potential Life Lost (YPLL) before age 75 by Cause of Death, Lancaster County, NE, 1999-2003**



- Lancaster County had a lower unintentional injury death rate (31.2 per 100,000), a higher suicide rate (11.1 per 100,000) and a lower homicide rate (2.0 per 100,000) than Nebraska rates (33.3, 10.6, and 3.2 per 100,000 respectively) (Table D3).

**Table D3: Injury Deaths, Percentage, and Age-adjusted Rates, Lancaster County, NE, 1999-2003**

Intent	Male			Female			Total			State		
	N	%	Rate	N	%	Rate	N	%	Rate	N	%	Rate
Unintentional	213	58.4	39.6	161	79.3	23.0	374	65.8	31.2	3,020	68.3	33.3
Suicide	116	31.8	18.9	21	10.3	3.4	137	24.1	11.1	894	20.2	10.6
Homicide	18	4.9	2.7	9	4.4	1.3	27	4.8	2.0	273	6.2	3.2
Undetermined	13	3.6	2.2	8	3.9	1.2	21	3.7	1.7	139	3.1	1.6
Adverse Effects	5	1.4	0.9	4	2.0	*	9	1.6	0.7	93	2.1	1.0
<b>Total</b>	<b>365</b>	<b>100.0</b>	<b>64.4</b>	<b>203</b>	<b>100.0</b>	<b>29.4</b>	<b>568</b>	<b>100.0</b>	<b>46.8</b>	<b>4,419</b>	<b>100.0</b>	<b>48.9</b>

Source: DHHS Death Certificate Data 1999-2003

\* Fewer than five deaths


- More males (64%) died from all injury-related causes than females (36%) (Table D4).
- Injury death rates were the highest for the oldest age group, 85 years and older. Persons in this age group had a higher injury death rate than the Nebraska rate (531 vs. 394 per 100,000 respectively) (Figure D2).

**Table D4: Injury-Related Deaths and Hospital Discharges by Gender, Age, and Race,  
Lancaster County, NE, 1999-2003**

	Lancaster County				State			
	Deaths		Hospital Discharges		Deaths		Hospital Discharges	
	N	%	N	%	N	%	N	%
<b>Gender</b>								
Male	365	64.3	58,734	53.5	2,848	64.3	386,563	53.1
Female	203	35.7	51,119	46.5	1,581	35.7	341,506	46.9
Total	568	100.0	109,853	100.0	4,429	100.0	728,069	100.0
<b>Age</b>								
<1	4	0.7	1,101	1.0	26	0.6	7,142	1.0
1-4	7	1.2	9,159	8.3	58	1.3	59,764	8.2
5-14	7	1.2	16,143	14.7	107	2.4	122,382	16.8
15-24	85	15.0	23,039	21.0	699	15.8	144,001	19.8
25-34	69	12.2	16,190	14.7	486	11.0	84,574	11.6
35-44	90	15.9	14,551	13.3	593	13.4	84,445	11.6
45-54	85	15.0	10,981	10.0	547	12.4	66,321	9.1
55-64	34	6.0	5,665	5.2	292	6.6	39,540	5.4
65-74	32	5.6	4,518	4.1	355	8.0	40,555	5.6
75-84	59	10.4	5,040	4.6	575	13.0	45,135	6.2
85+	96	16.9	3,469	3.2	691	15.6	34,222	4.7
Total	568	100.0	109,856	100.0	4,429	100.0	728,081	100.0
<b>Race</b>								
Asian	8	1.4			29	0.7		
African American	13	2.3			194	4.4		
Native American	14	2.5			72	1.6		
White	532	93.7			4,123	93.1		
Other	1	0.2			11	0.3		
Total	568	100.0			4,429	100.0		

Note: Hospital discharges include hospital inpatients, outpatients, and ER visits.

**Figure D2: All Injury Death Rates by Age, Lancaster County, NE, 1999-2003**




\* Fewer than five deaths  
Source: DHHS Death Certificate Data 1999-2003

**Leading causes**

- Suicide was the leading cause of injury deaths with an age-adjusted rate of 11.1 per 100,000. The rate was slightly higher than the Nebraska rate (10.6 per 100,000) (Table D3 & Figure D3).
- Motor vehicle crashes were the second leading cause of injury deaths. The age-adjusted death rate for motor vehicle crashes was lower than the Nebraska rate (10.0 vs. 13.6 per 100,000, respectively) (Figure D3). Lancaster County has not met the 2010 Nebraska Objective of 12.0 or fewer motor vehicle crash injury deaths per 100,000 population.
- Falls were the third leading cause of injury deaths. The age-adjusted death rate due to falls (8.8 per 100,000) was slightly higher than the Nebraska rate (7.0 per 100,000) (Figure D3). Lancaster County has not met the 2010 Nebraska Objective of 3.5 or fewer fall injury deaths per 100,000 population.

**Figure D3: Age-adjusted Injury Death Rates by Cause, Lancaster County, NE, 1999-2003**



Source: DHHS Death Certificate Data 1999-2003

**Age**

- The most motor vehicle crash deaths were among persons age 15-24 years (N=51), accounting for 38.3% of total deaths due to motor vehicle crashes (N=133) (Table D5).

**Table D5: Deaths Due to Motor Vehicle Crash by Age and Gender, Lancaster County, 1999-2003**

Age	Male		Female		Total	
	N	%	N	%	N	%
1-4	1	1.2	0	0.0	1	0.8
5-14	3	3.5	1	2.1	4	3.0
15-24	33	38.4	18	38.3	51	38.3
25-34	13	15.1	6	12.8	19	14.3
35-44	13	15.1	3	6.4	16	12.0
45-54	11	12.8	5	10.6	16	12.0
55-64	3	3.5	4	8.5	7	5.3
65-74	1	1.2	3	6.4	4	3.0
75-84	5	5.8	4	8.5	9	6.8
85+	3	3.5	3	6.4	6	4.5
<b>Total</b>	<b>86</b>	<b>100.0</b>	<b>47</b>	<b>100.0</b>	<b>133</b>	<b>100.0</b>

Source: DHHS Death Certificate Data 1999-2003

- Persons 85 years and older (N=53) accounted for over half (57%) of deaths due to fall injuries (Table D6).

**Table D6: Deaths Due to Fall by Age and Gender,  
Lancaster County, 1999-2003**

Age	Male		Female		Total	
	N	%	N	%	N	%
1-4	1	2.4	0	0.0	1	1.1
25-34	1	2.4	0	0.0	1	1.1
45-54	2	4.9	0	0.0	2	2.2
55-64	6	14.6	0	0.0	6	6.5
65-74	3	7.3	3	5.8	6	6.5
75-84	8	19.5	16	30.8	24	25.8
85+	20	48.8	33	63.5	53	57.0
<b>Total</b>	<b>41</b>	<b>100.0</b>	<b>52</b>	<b>100.0</b>	<b>93</b>	<b>100.0</b>

Source: DHHS Death Certificate Data 1999-2003

- Persons in the 35-44 year age group had the most deaths (N=38) due to suicide (*Table D7*).

**Table D7: Deaths Due to Suicide by Age and Gender,  
Lancaster County, 1999-2003**

Age	Male		Female		Total	
	N	%	N	%	N	%
5-14	1	0.9	0	0.0	1	0.7
15-24	12	10.4	2	9.5	14	10.3
25-34	26	22.6	4	19.0	30	22.1
35-44	31	27.0	7	33.3	38	27.9
45-54	24	20.9	5	23.8	29	21.3
55-64	8	7.0	2	9.5	10	7.4
65-74	8	7.0	0	0.0	8	5.9
75-84	5	4.3	1	4.8	6	4.4
85+	1	0.9	0	0.0	1	0.7
<b>Total</b>	<b>115</b>	<b>100.0</b>	<b>21</b>	<b>100.0</b>	<b>136</b>	<b>100.0</b>

Source: DHHS Death Certificate Data 1999-2003

**Gender**

- More males (N=86) died from motor vehicle crashes than females (N=47) (*Table D5*).
- More females (N=52) died from falls than males (N=41) (*Table D6*).
- The number of males (N=115) who died from suicide was more than five times higher than females (N=21) (*Table D7*).

## Hospital discharges

### Overall

- Lancaster County had a higher unintentional injury hospital discharge rate (8,232 per 100,000), a lower suicide attempt rate (66 per 100,000) and assault rate (197 per 100,000) than Nebraska rates (7,610, 75, and 204 per 100,000 respectively) (*Table H1*).
- More males (N=58,092) than females (N=50,474) were treated at a hospital for all injury-related causes (*Table H1*).

**Table H1: Injury Hospital Discharges, Percentage, and Age-adjusted Rates, Lancaster County, NE, 1999-2003**

Intent	Males			Females			Total			State		
	N	%	Rate	N	%	Rate	N	%	Rate	N	%	Rate
Unintentional	55,881	96.2	8,816.9	48,766	96.6	7,652.8	104,647	96.4	8,232.4	665,407	95.9	7,609.6
Suicide	328	0.6	47.1	568	1.1	83.8	896	0.8	65.5	6,379	0.9	75.0
Homicide	1,743	3.0	239.6	1,056	2.1	153.6	2,799	2.6	196.5	17,720	2.6	203.9
Undetermined	61	0.1	9.4	59	0.1	8.8	120	0.1	9.1	4,181	0.6	48.3
Other	79	0.1	11.3	25	0.0	3.6	104	0.1	7.5	404	0.1	4.7
<b>Total</b>	<b>58,092</b>	<b>100.0</b>	<b>9,124.3</b>	<b>50,474</b>	<b>100.0</b>	<b>7,902.7</b>	<b>108,566</b>	<b>100.0</b>	<b>8,511.0</b>	<b>694,091</b>	<b>100.0</b>	<b>7,941.5</b>

Source: Nebraska Hospital Discharge Data 1999-2003

Rates per 100,000 population are age-adjusted to the 2000 U.S. standard population

### Leading causes

- Falls were the leading cause of injury hospital discharge for all age groups combined, with an age-adjusted rate of 2,384 per 100,000. The rate was relatively higher than the Nebraska rate (2,184 per 100,000). Falls were also the leading cause of hospital discharge for all age groups except for the 10-14 and 15-24 year age groups (*Table H2 & Figure H1*).
- Struck by/against was the second leading cause of injury hospital discharge for all age groups combined, and the leading cause for persons age 10-14. The age-adjusted hospital discharge rate for struck by/against was slightly higher than the state rate (1,154 vs. 1,132 per 100,000 respectively) (*Table H2 & Figure H1*).
- Motor vehicle crashes were the third leading cause of injury hospital discharge for all age groups combined, and the leading cause for persons age 15-24. The age-adjusted rate was higher than the Nebraska rate (869 vs. 684 per 100,000) (*Table H2 & Figure H1*).
- Overexertion (808 per 100,000) and cutting/piercing (674 per 100,000) were the 4<sup>th</sup> and 5<sup>th</sup> ranking causes of injury hospital discharges in Lancaster County (*Table H2 & Figure H1*).

**Table H2: Five Leading Causes of Hospital Discharges for Injury by Age, Lancaster County, NE, 1999-2003**


Rank	<1	1-4	5-9	10-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	84+	All Ages
1	Falls N=411	Falls N=3379	Falls N=2405	Struck by/against N=2357	MV traffic N=4156	Falls N=2457	Falls N=2702	Falls N=2687	Falls N=1814	Falls N=2016	Falls N=3059	Falls N=2576	Falls N=28,923
2	Struck by/against N=152	Struck by/against N=1584	Struck by/against N=1525	Falls N=2327	Struck by/against N=3983	MV traffic N=2452	Overexertion N=1934	MV traffic N=1259	Overexertion N=525	Overexertion N=326	Struck by/against N=238	Struck by/against N=110	Struck by/against N=14,963
3	Fire/Burn N=145	Fire/Burn N=705	Cut/Pierce N=598	Overexertion N=801	Falls N=3090	Overexertion N=2191	MV traffic N=1806	Overexertion N=1213	MV traffic N=513	MV traffic N=303	MV traffic N=235	Overexertion N=84	MV traffic N=11,765
4	Cut/Pierce N=40	Cut/Pierce N=456	Other Pedal CYC N=480	Cut/Pierce N=628	Overexertion N=2635	Struck by/against N=1944	Struck by/against N=1525	Struck by/against N=945	Cut/Pierce N=477	Cut/Pierce N=294	Overexertion N=230	MV traffic N=80	Overexertion N=10,612
5	Natural/Envir N=31	Overexertion N=441	MV traffic N=311	Other Pedal CYC N=575	Cut/Pierce N=2321	Cut/Pierce N=1623	Cut/Pierce N=1269	Cut/Pierce N=916	Struck by/against N=358	Struck by/against N=243	Cut/Pierce N=152	Cut/Pierce N=38	Cut/Pierce N=8811

Hospital visits includes visit as inpatient, ER and non-ER patient

Note: Causes coded as "other specified" and "unspecified" are excluded in this matrix

Source: Nebraska Hospital Discharge Data 1999-2003

**Figure H1: Age-adjusted Injury Hospital Discharge Rates by Cause, Lancaster County, NE, 1999-2003**



Source: Nebraska Hospital Discharge Data 1999-2003

### Age

- The hospital discharge rate for falls was the highest for persons age 85 and older. Among children age 14 and under, the 1-4 year age group had the highest fall injury hospital discharge rate (5,031 per 100,000), which was relatively higher than the Nebraska rate (4,310 per 100,000). This age group had the third highest fall injury hospital discharge rate after the 75-84 and 85 and older age groups (*Table H3*).
- The 5-14 year age group had the highest hospital discharge rate (2,434 per 100,000) for struck by/against, which was higher than the Nebraska rate (2,237 per 100,000). The rate was second highest for the 1-4 year age group at 2,359 per 100,000 (*Table H3*).
- The hospital discharge rate for motor vehicle crashes was highest for the 15-24 year age group (1,753 per 100,000) (*Table H3*).
- The hospital discharge rate for overexertion injuries was the highest for individuals in the 25-34 year age group (1,119 per 100,000). The hospital discharge rate for cutting/piercing injuries were the highest for individuals in the 15-24 year age group (979 per 100,000) (*Table H3*).

### Gender

- Females had higher injury hospital discharge rates for falls than males (2,561 vs. 2,204 per 100,000) (*Table H3*).
- The injury hospital discharge rate for struck by/against for males was almost twice that for females (1,474 vs. 836 per 100,000) (*Table H3*).

- Females had higher hospital discharge rate for motor vehicle crashes than males (963 vs. 774 per 100,000) (*Table H3*).

**Table H3: Five Leading Causes of Injury Hospital Discharges, Rates by Gender and Age, Lancaster County, NE, 1999-2003**

Cause	Age	Lancaster County									State		
		Male			Female			Total			Rate		
		N	%	Rate	N	%	Rate	N	%	Rate	Male	Female	Total
Falls	<1	216	1.7	2329.6	195	1.2	2181.9	411	1.4	2257.1	2298.5	2139.4	2220.8
Falls	1-4	1998	15.7	5872.3	1381	8.5	4167.9	3379	11.7	5031.4	4871.2	3721.2	4309.7
Falls	5-14	2663	20.9	3258.5	2069	12.8	2662	4732	16.4	2967.7	2855.0	2404.1	2634.8
Falls	15-24	1683	13.2	1400.2	1407	8.7	1203.7	3090	10.7	1303.4	1570.9	1323.5	1449.9
Falls	25-34	1194	9.4	1160.9	1263	7.8	1358.6	2457	8.5	1254.8	1048.4	1303.4	1173.8
Falls	35-44	1235	9.7	1286.8	1467	9.1	1601.8	2702	9.3	1440.6	1084.2	1357.6	1220.2
Falls	45-54	1062	8.3	1285.3	1625	10	1936.7	2687	9.3	1613.5	1116.0	1610.3	1364.2
Falls	55-64	698	5.5	1506.1	1116	6.9	2287.8	1814	6.3	1906.9	1337.7	1976.8	1664.3
Falls	65-74	642	5	2146.2	1374	8.5	3770.3	2016	7	3038.2	2151.5	3581.0	2926.8
Falls	75-84	854	6.7	4541.3	2205	13.6	7782.7	3059	10.6	6489.6	4355.2	7355.0	6138.2
Falls	85+	508	4	10453	2068	12.8	15651	2576	8.9	14253	10070.0	14640.0	13320.0
<b>Total</b>		12753	100.0	2204.3	16170	100.0	2561.2	28923	100.0	2384.0	2013.8	2349.3	2184.0
Struck by/against	<1	75	0.8	808.9	77	1.5	861.6	152	1	834.8	836	692.9	766.1
Struck by/against	1-4	995	10.3	2924.4	589	11.2	1777.6	1584	10.6	2358.6	2812.4	1729.5	2283.7
Struck by/against	5-14	2591	26.7	3170.4	1291	24.5	1661	3882	25.9	2434.6	2919.3	1521.6	2236.7
Struck by/against	15-24	2837	29.2	2360.4	1145	21.8	979.6	3982	26.6	1679.6	3069.3	1161.5	2135.8
Struck by/against	25-34	1268	13.1	1232.9	676	12.8	727.2	1944	13	992.8	1212	676.9	948.9
Struck by/against	35-44	967	10	1007.6	558	10.6	609.3	1525	10.2	813.1	854.1	527.3	691.6
Struck by/against	45-54	542	5.6	655.9	403	7.7	480.3	945	6.3	567.5	574.9	385.7	479.9
Struck by/against	55-64	186	1.9	401.3	172	3.3	352.6	358	2.4	376.3	414.7	291.2	351.6
Struck by/against	65-74	114	1.2	381.1	129	2.5	354	243	1.6	366.2	415.7	292.7	349
Struck by/against	75-84	96	1	510.5	142	2.7	501.2	238	1.6	504.9	415.6	363.7	384.7
Struck by/against	85+	31	0.3	637.9	79	1.5	597.9	110	0.7	608.6	507.4	519.4	516
<b>Total</b>		9702	100.0	1474.1	5261	100.0	836.1	14963	100.0	1154.0	1488.5	786.4	1132.3
MV traffic	<1	14	0.3	151	11	0.2	123.1	25	0.2	137.3	128.9	121.4	125.2
MV traffic	1-4	78	1.5	229.2	81	1.2	244.5	159	1.3	236.8	251.0	229.3	240.4
MV traffic	5-14	389	7.4	476	418	6.4	537.8	807	6.8	506.1	400.0	436.0	417.6
MV traffic	15-24	1761	33.4	1465.1	2395	36.7	2049	4156	35.2	1753	1444.2	1831.5	1633.7
MV traffic	25-34	1172	22.2	1139.5	1280	19.6	1376.9	2452	20.8	1252.2	855.6	993.0	923.2
MV traffic	35-44	824	15.6	858.6	982	15.1	1072.2	1806	15.3	962.9	607.3	708.3	657.6
MV traffic	45-54	579	11	700.7	680	10.4	810.4	1259	10.7	756	476.5	523.1	499.9
MV traffic	55-64	217	4.1	468.2	296	4.5	606.8	513	4.3	539.3	346.5	441.2	394.9
MV traffic	65-74	120	2.3	401.2	183	2.8	502.2	303	2.6	456.6	327.3	382.2	357.1
MV traffic	75-84	92	1.7	489.2	143	2.2	504.7	235	2	498.5	402.4	393.8	397.3
MV traffic	85+	28	0.5	576.1	52	0.8	393.6	80	0.7	442.6	404.8	248.5	293.6
<b>Total</b>		5274	100.0	774.0	6521	100.0	962.7	11795	100.0	869.0	629.3	737.3	684.0
Overexertion	0-1	13	0.2	140.2	16	0.3	179	29	0.3	159.3	111.1	156.8	133.4
Overexertion	1-4	211	3.7	620.2	230	4.7	694.2	441	4.2	656.7	352.6	519.6	434.1
Overexertion	5-14	499	8.8	610.6	506	10.2	651	1005	9.5	630.3	537.1	610.3	572.9
Overexertion	15-24	1529	27	1272.1	1105	22.4	945.4	2634	24.8	1111	1317.5	1024.1	1173.9
Overexertion	25-34	1252	22.1	1217.3	939	19	1010.1	2191	20.6	1118.9	952.4	828.7	891.6
Overexertion	35-44	1084	19.1	1129.5	850	17.2	928.1	1934	18.2	1031.2	817.7	677.3	747.9
Overexertion	45-54	619	10.9	749.1	594	12	707.9	1213	11.4	728.4	616.4	572.9	594.5
Overexertion	55-64	241	4.2	520	284	5.8	582.2	525	4.9	551.9	426.6	433.8	430.3
Overexertion	65-74	132	2.3	441.3	194	3.9	532.3	326	3.1	491.3	356.1	445.4	404.6
Overexertion	75-84	78	1.4	414.8	152	3.1	536.5	230	2.2	487.9	311.8	419.9	376
Overexertion	85+	15	0.3	308.6	69	1.4	522.2	84	0.8	464.8	236.9	387.2	343.8
<b>Total</b>		5673	100.0	849.1	4939	100.0	768.1	10612	100.0	808.0	705.5	635.6	679.2
Cut/Pierce	<1	24	0.4	258.8	16	0.5	179	40	0.5	219.7	228.7	185.5	207.6
Cut/Pierce	1-4	271	4.7	796.5	185	6	558.3	456	5.2	679	925.2	609.5	771.1
Cut/Pierce	5-14	783	13.7	958.1	442	14.3	568.7	1225	13.9	768.3	1033.2	615.8	829.4
Cut/Pierce	15-24	1591	27.8	1323.7	730	23.6	624.5	2321	26.3	979	1399.6	650.1	1032.9
Cut/Pierce	25-34	1107	19.4	1076.3	516	16.7	555.1	1623	18.4	828.9	1138.4	596.8	872.2
Cut/Pierce	35-44	796	13.9	829.4	473	15.3	516.5	1269	14.4	676.6	861.0	517.5	690.1
Cut/Pierce	45-54	567	9.9	686.2	349	11.3	415.9	916	10.4	550	650.7	401.2	525.4
Cut/Pierce	55-64	301	5.3	649.5	176	5.7	360.8	477	5.4	501.4	555.0	309.7	429.6
Cut/Pierce	65-74	172	3	575	122	3.9	334.8	294	3.3	443.1	582.4	266.5	411.1
Cut/Pierce	75-84	86	1.5	457.3	66	2.1	233	152	1.7	322.5	448.3	214.8	309.5
Cut/Pierce	85+	17	0.3	349.8	21	0.7	158.9	38	0.4	210.3	333.7	194.8	234.9
<b>Total</b>		5715	100.0	864.5	3096	100.0	484.5	8811	100.0	674.0	893.0	492.5	689.8

Source: Nebraska Hospital Discharge Data 1999-2003  
Rates are per 100,000 per year