

Table 1C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Demographic Indicators ⁽²⁾

Total Records	Number	%	Source of Data	Number	%
Complete Records	7,594	73.8	WIC	10,293	100.0
Prenatal Records	694	6.7	Maternal Health	0	0.0
Postpartum Records	2,005	19.5	Other	0	0.0
Total Records Accepted	10,293		Total	10,293	100.0
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%
White, Not Hispanic	5,796	56.3
Black, Not Hispanic	1,183	11.5
Hispanic	2,779	27.0
American Indian/Alaska Native	149	1.4
Asian/Pacific Islander	270	2.6
Multiple Races	115	1.1
All Other	0	0.0
Total	10,292	100.0
<i>Excluded Unknown: 1</i>	<i>0.0</i>	

Age Distribution	Number	%
< 15 Years	21	0.2
15 - 17 Years	468	4.5
18 - 19 Years	1,164	11.3
20 - 29 Years	6,341	61.6
30 - 39 Years	2,146	20.9
>= 40 Years	146	1.4
Total	10,286	100.0
<i>Excluded Unknown: 7</i>	<i>0.1</i>	

Education Distribution	Number	%
< High School	3,026	29.5
High School	4,236	41.3
> High School	2,999	29.2
Total	10,261	100.0
<i>Excluded Unknown: 32</i>	<i>0.3</i>	

% Poverty Level	Number	%
0 - 50	3,006	36.3
51 - 100	2,767	33.4
101 - 130	1,246	15.0
131 - 150	543	6.6
151 - 185	593	7.2
186 - 200	49	0.6
> 200	79	1.0
Adjunctive Eligibility	0	0.0
Total	8,283	100.0
<i>Excluded Unknown: 5</i>	<i>0.1</i>	

Program Participation ⁽³⁾	Number	%
WIC	8,275	100.0
SNAP	506	30.2
Medicaid	506	66.4
TANF	506	7.5
<i>Excluded Unknown:</i>		
<i>WIC:</i>	13	0.2
<i>SNAP:</i>	7,782	93.9
<i>Medicaid:</i>	7,782	93.9
<i>TANF:</i>	7,782	93.9

Migrant Status	Number	%
Yes	7	0.1
No	10,269	99.9
Total	10,276	100.0
<i>Excluded Unknown: 17</i>	<i>0.2</i>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Health Indicators

Summary of Health Indicators (2)				
	Nebraska Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	9,975	3.7	1,106,370	4.5
Normal Weight	9,975	40.0	1,106,370	42.1
Overweight	9,975	26.2	1,106,370	26.0
Obese	9,975	30.1	1,106,370	27.4
Maternal Weight Gain (4)				
Less Than Ideal	9,236	22.2	835,927	21.5
Ideal	9,236	31.2	835,927	30.6
Greater Than Ideal	9,236	46.6	835,927	48.0
Anemia (Low Hb/Hct) (5)				
1st Trimester	2,695	6.9	419,016	7.6
2nd Trimester	3,122	7.9	333,582	12.0
3rd Trimester	1,219	23.9	164,201	33.9
Postpartum	2,708	19.6	346,981	28.1
Parity (6)				
0	10,251	30.4	965,206	40.3
1	10,251	25.5	965,206	26.8
2	10,251	18.2	965,206	16.7
≥ 3	10,251	25.8	965,206	16.3
Interpregnancy Interval				
< 6 Months	1,769	3.7	554,361	11.7
6 - <18 Months	1,769	8.6	554,361	25.9
≥18 Months	1,769	87.7	554,361	62.4
Gestational Diabetes	0	*	419,956	5.2
Hypertension During Pregnancy	28	*	467,675	6.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Health Indicators

Summary of Health Indicators (2)				
	Nebraska Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	9,576	85.7	429,770	83.1
No Care	9,576	4.5	429,770	4.1
WIC Enrollment				
1st Trimester	10,018	31.2	1,074,863	36.0
2nd Trimester	10,018	34.2	1,074,863	33.9
3rd Trimester	10,018	15.0	1,074,863	16.9
Postpartum (3)	10,018	19.5	1,074,863	13.2
Prepregnancy Multivitamin Consumption (4)	0	*	380,154	19.2
Multivitamin Consumption During Pregnancy (5)	4	*	418,448	85.0
Smoking				
3 Months Prior to Pregnancy	10,263	31.7	814,631	23.0
During Pregnancy	7,909	16.2	710,227	14.0
Last 3 Months of Pregnancy	9,558	14.7	715,034	12.1
Postpartum	9,591	16.6	722,233	14.1
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	2,520	49.7	166,084	43.5
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	2,245	44.0	137,883	38.1
Smoking in Household				
During Pregnancy	107	100.0	660,570	16.3
Postpartum	1,113	100.0	562,126	15.6
Drinking				
3 Months Prior to Pregnancy	10,227	18.9	930,233	8.8
Last 3 Months of Pregnancy	9,516	0.2	577,187	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Health Indicators

Summary of Health Indicators (2)				
	Nebraska Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	9,223	0.9	880,933	1.1
Low	9,223	6.4	880,933	6.9
Normal	9,223	85.6	880,933	85.3
High	9,223	7.0	880,933	6.7
Preterm (4)	9,151	9.7	948,286	10.7
Full term Low Birth Weight (5)	8,232	2.9	749,107	3.4
Multiple Births	9,451	1.3	978,390	1.4
Breastfeeding				
Ever Breastfed	9,261	75.6	871,811	68.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.



Table 3C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Comparison of Racial and Ethnic Distribution by Contributor to Nation

Comparison of Racial and Ethnic Distribution ⁽²⁾								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Nebraska	10,293	56.3	11.5	27.0	1.4	2.6	1.1	0.0
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Comparison of Age Distribution by Contributor to Nation

Comparison of Age Distribution ⁽²⁾								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Nebraska	10,293	0.2	4.5	11.3	61.6	20.8	1.4	0.1
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Comparison of Maternal Health Indicators by Contributor to Nation

Comparison of Maternal Health Indicators ⁽²⁾														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
		Number	%		%	Number	%	%	Number	%			Number	%
Nebraska	9,975	3.7	56.3	9,236	22.2	46.6	1,219	23.9	2,708	19.6	0	*	28	*
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Comparison of Maternal Behavioral Indicators by Contributor to Nation

Comparison of Maternal Behavioral Indicators ⁽²⁾									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Nebraska	9,576	85.7	4.5	10,018	31.2	0	*	4	*
Nation (Prior Year)	429,770	83.1	4.1	1,074,863	36.0	380,154	19.2	418,448	85.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Comparison of Smoking Indicators by Contributor to Nation

Comparison of Smoking Indicators (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	%	Number	%	Number	%	Number	%		
Nebraska	10,263	31.7	9,558	14.7	2,245	44.0	107	100.0	1,113	100.0
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Comparison of Infant Health Indicators by Contributor to Nation

Comparison of Infant Health Indicators ⁽²⁾									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
Nebraska	9,223	7.4	7.0	9,151	9.7	8,232	2.9	9,261	75.6
Nation (Prior Year)	880,933	8.0	6.7	948,286	10.7	749,107	3.4	871,811	68.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 9C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Maternal Health Indicators by Race/Ethnicity, Age, or Education

Maternal Health Indicators by Race/Ethnicity, Age, or Education ⁽²⁾														
	Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾				Gestational Diabetes		Hypertension During Pregnancy	
	Number	Under-weight	Over-weight	Number	< Ideal	> Ideal	3rd Trimester		Postpartum		Number	%	Number	%
		%	%		%	%	Number	%	Number	%				
Race/Ethnicity														
White, Not Hispanic	5,636	4.1	55.2	5,215	19.7	50.7	666	21.0	1,611	14.3	0	*	19	*
Black, Not Hispanic	1,153	4.4	56.3	1,044	25.6	43.7	127	47.2	416	35.8	0	*	2	*
Hispanic	2,674	1.9	61.2	2,521	25.4	40.7	360	20.3	520	21.9	0	*	5	*
American Indian/Alaska Native	142	2.1	64.1	117	22.2	49.6	23	*	34	*	0	*	1	*
Asian/Pacific Islander	256	12.5	26.2	240	33.8	26.7	31	*	106	24.5	0	*	0	*
Multiple Races	113	3.5	54.9	98	*	*	12	*	20	*	0	*	1	*
All Other/Unknown	1	*	*	1	*	*	0	*	1	*	0	*	0	*
Total	9,975	3.7	56.3	9,236	22.2	46.6	1,219	23.9	2,708	19.6	0	*	28	*
Age														
< 15 Years	20	*	*	18	*	*	3	*	6	*	0	*	0	*
15 - 17 Years	456	4.8	35.7	418	19.9	50.0	66	*	107	24.3	0	*	1	*
18 - 19 Years	1,122	7.0	41.7	1,020	17.8	51.4	139	22.3	286	24.8	0	*	2	*
20 - 29 Years	6,170	3.8	55.4	5,712	22.1	46.7	778	23.7	1,700	18.9	0	*	17	*
30 - 39 Years	2,062	1.8	70.1	1,932	25.0	43.6	220	23.2	570	17.9	0	*	8	*
>= 40 Years	138	0.7	78.3	129	28.7	36.4	13	*	39	*	0	*	0	*
Unknown	7	*	*	7	*	*	0	*	0	*	0	*	0	*
Total	9,975	3.7	56.3	9,236	22.2	46.6	1,219	23.9	2,708	19.6	0	*	28	*
Education														
< High School	2,930	4.0	55.8	2,706	26.5	41.6	382	26.7	683	24.5	0	*	8	*
High School	4,121	3.8	55.3	3,818	20.8	48.6	510	22.2	1,115	19.8	0	*	14	*
> High School	2,919	3.3	58.2	2,708	19.8	48.8	327	23.2	906	15.6	0	*	6	*
Unknown	5	*	*	4	*	*	0	*	4	*	0	*	0	*
Total	9,975	3.7	56.3	9,236	22.2	46.6	1,219	23.9	2,708	19.6	0	*	28	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 10C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education (2)									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	5,385	87.9	4.3	5,644	32.0	0	*	3	*
Black, Not Hispanic	1,074	86.9	2.6	1,161	28.5	0	*	0	*
Hispanic	2,635	81.1	6.0	2,687	31.6	0	*	0	*
American Indian/Alaska Native	122	81.1	4.9	146	28.1	0	*	1	*
Asian/Pacific Islander	257	84.0	3.1	268	26.1	0	*	0	*
Multiple Races	102	82.4	4.9	111	26.1	0	*	0	*
All Other/Unknown	1	*	*	1	*	0	*	0	*
Total	9,576	85.7	4.5	10,018	31.2	0	*	4	*
Age									
< 15 Years	20	*	*	21	*	0	*	0	*
15 - 17 Years	431	77.0	5.8	460	34.8	0	*	1	*
18 - 19 Years	1,066	82.2	5.5	1,145	35.5	0	*	0	*
20 - 29 Years	5,894	87.0	4.2	6,181	30.4	0	*	3	*
30 - 39 Years	2,022	86.1	4.5	2,067	31.1	0	*	0	*
>= 40 Years	136	84.6	7.4	137	23.4	0	*	0	*
Unknown	7	*	*	7	*	0	*	0	*
Total	9,576	85.7	4.5	10,018	31.2	0	*	4	*
Education									
< High School	2,821	80.0	5.8	2,922	31.3	0	*	2	*
High School	3,951	86.5	4.4	4,125	33.1	0	*	2	*
> High School	2,800	90.3	3.4	2,942	28.5	0	*	0	*
Unknown	4	*	*	29	*	0	*	0	*
Total	9,576	85.7	4.5	10,018	31.2	0	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 11C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Smoking Indicators by Race/Ethnicity, Age, or Education

Smoking Indicators by Race/Ethnicity, Age, or Education ⁽²⁾										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	5,776	44.9	5,375	22.5	1,782	40.1	68	*	884	100.0
Black, Not Hispanic	1,180	22.9	1,072	9.3	185	49.7	16	*	83	*
Hispanic	2,774	8.9	2,631	1.9	180	70.6	9	*	107	100.0
American Indian/Alaska Native	148	43.9	121	22.3	46	*	7	*	11	*
Asian/Pacific Islander	269	7.4	256	0.8	12	*	3	*	7	*
Multiple Races	115	48.7	102	17.6	40	*	4	*	21	*
All Other/Unknown	1	*	1	*	0	*	0	*	0	*
Total	10,263	31.7	9,558	14.7	2,245	44.0	107	100.0	1,113	100.0
Age										
< 15 Years	21	*	20	*	0	*	0	*	3	*
15 - 17 Years	467	22.3	430	7.0	84	*	6	*	52	*
18 - 19 Years	1,161	36.2	1,066	12.2	311	55.0	14	*	146	100.0
20 - 29 Years	6,319	35.1	5,878	16.7	1,520	42.3	69	*	736	100.0
30 - 39 Years	2,142	23.1	2,020	12.7	320	35.9	17	*	169	100.0
>= 40 Years	146	14.4	137	5.8	10	*	1	*	7	*
Unknown	7	*	7	*	0	*	0	*	0	*
Total	10,263	31.7	9,558	14.7	2,245	44.0	107	100.0	1,113	100.0
Education										
< High School	3,025	26.3	2,816	13.8	560	40.9	41	*	291	100.0
High School	4,235	37.1	3,941	17.7	1,102	41.1	43	*	564	100.0
> High School	2,999	29.6	2,797	11.5	583	52.3	23	*	257	100.0
Unknown	4	*	4	*	0	*	0	*	1	*
Total	10,263	31.7	9,558	14.7	2,245	44.0	107	100.0	1,113	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 12C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Infant Health Indicators by Race/Ethnicity, Age, or Education

Infant Health Indicators by Race/Ethnicity, Age, or Education (2)									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	5,209	6.9	7.4	5,169	9.8	4,647	2.4	5,226	73.7
Black, Not Hispanic	1,042	11.7	4.3	1,037	11.0	919	5.0	1,047	70.1
Hispanic	2,508	6.5	7.8	2,483	9.3	2,242	2.7	2,522	82.4
American Indian/Alaska Native	118	4.2	10.2	116	6.9	108	3.7	118	65.3
Asian/Pacific Islander	244	10.2	2.9	245	9.0	221	5.9	246	76.8
Multiple Races	101	5.9	4.0	100	6.0	94	*	101	69.3
All Other/Unknown	1	*	*	1	*	1	*	1	*
Total	9,223	7.4	7.0	9,151	9.7	8,232	2.9	9,261	75.6
Age									
< 15 Years	19	*	*	19	*	17	*	19	*
15 - 17 Years	421	7.8	5.2	420	8.6	382	3.7	423	73.0
18 - 19 Years	1,041	4.6	4.8	1,036	8.9	944	2.2	1,042	74.6
20 - 29 Years	5,706	6.4	6.7	5,661	8.9	5,137	2.8	5,725	75.2
30 - 39 Years	1,907	7.4	9.8	1,889	12.7	1,641	3.2	1,921	77.8
>= 40 Years	122	5.7	5.7	119	10.9	105	2.9	124	77.4
Unknown	7	*	*	7	*	6	*	7	*
Total	9,223	7.4	7.0	9,151	9.7	8,232	2.9	9,261	75.6
Education									
< High School	2,702	7.0	6.7	2,667	9.7	2,399	3.7	2,715	73.5
High School	3,803	6.5	6.6	3,779	9.9	3,392	2.7	3,819	72.9
> High School	2,712	5.8	8.0	2,702	9.5	2,438	2.4	2,722	81.4
Unknown	6	*	*	3	*	3	*	5	*
Total	9,223	7.4	7.0	9,151	9.7	8,232	2.9	9,261	75.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 13C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska

Maternal Weight Gain and Birth Outcomes by Select Health Indicators

Maternal Weight Gain and Birth Outcomes by Select Health Indicators (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal	> Ideal		Birthweight			Preterm (6)	Full Term LBW (7)		
		Number	%	%	Low (4)	High (5)				
	Number	%	%	Number	%	%	Number	%	Number	%
Prepregnancy BMI										
Underweight	329	31.6	29.5	327	10.4	1.8	326	9.2	293	4.8
Normal Weight	3,694	26.3	36.6	3,616	7.4	4.9	3,600	9.7	3,236	3.2
Overweight	2,430	14.3	56.0	2,363	7.4	6.6	2,349	9.6	2,117	2.6
Obese	2,783	22.6	53.6	2,697	6.7	10.6	2,662	9.8	2,398	2.3
Weight Gain										
< Ideal	-	-	-	1,888	11.3	3.2	1,869	13.3	1,616	4.4
Ideal	-	-	-	2,827	7.4	5.4	2,811	9.9	2,519	3.5
> Ideal	-	-	-	4,242	5.5	9.7	4,215	7.9	3,869	1.8
Smoking During Pregnancy										
Smokers	1,097	20.7	52.2	1,102	9.1	4.5	1,090	9.8	979	3.9
Non-Smokers	5,841	21.9	46.1	5,871	6.3	7.5	5,837	9.2	5,282	2.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Racial and Ethnic Distribution

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2011	10,293	56.3	11.5	27.0	1.4	2.6	1.1	0.0
2010	10,821	56.2	11.2	27.7	1.4	2.2	1.2	0.0
2009	11,420	55.1	10.9	29.7	1.3	1.9	1.1	0.0
2008	11,549	56.9	10.7	27.9	1.5	1.8	1.1	0.0
2007	11,858	57.0	10.8	27.9	1.8	1.7	0.9	0.0
2006	12,457	57.2	10.8	27.8	1.8	1.7	0.6	0.0
2005	11,821	57.5	9.9	28.1	1.9	2.1	0.6	0.1
2004	11,708	58.3	9.7	27.2	1.9	2.1	0.2	0.5
2003	11,814	59.5	9.0	26.8	1.6	2.0	0.0	1.1
2002	11,401	60.9	9.4	25.4	1.8	1.7	0.0	0.9
2001	11,016	62.9	9.7	23.2	1.5	1.9	0.0	0.8
2000	10,381	64.6	10.0	21.2	1.4	2.1	0.0	0.7
1999	9,737	66.0	10.4	19.4	1.3	1.9	0.0	1.0
1998	8,945	68.8	9.0	18.3	1.2	1.9	0.0	0.8
1997	4,132	73.5	5.5	17.8	1.0	1.5	0.0	0.7
1996	7,004	71.6	9.9	14.9	1.1	1.8	0.0	0.7
1995	0	*	*	*	*	*	*	*
1994	7,538	74.6	11.3	11.1	1.2	1.6	0.0	0.2
1993	5,955	76.9	10.7	9.4	1.3	1.6	0.0	0.1
1992	6,190	75.9	12.1	8.6	1.8	1.5	0.0	0.1
1991	6,208	76.1	12.8	7.4	2.2	1.4	0.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Racial and Ethnic Distribution

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1990	2,259	79.6	11.2	6.0	1.6	1.4	0.0	0.0
1989	2,452	78.1	12.7	5.5	1.9	1.3	0.0	0.6
1988	3,298	76.1	14.6	4.9	2.5	1.6	0.0	0.3
1987	3,790	75.5	16.2	4.7	2.1	1.3	0.0	0.2
1986	2,421	77.9	14.5	4.2	1.7	1.1	0.0	0.6
1985	724	78.9	10.9	2.6	1.5	5.7	0.0	0.4
1984	2,437	71.3	18.8	4.6	2.2	2.1	0.0	1.1
1983	2,492	71.8	18.9	4.8	2.0	1.6	0.0	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2011	10,293	0.2	4.5	11.3	61.6	20.8	1.4	0.1
2010	10,821	0.2	4.9	12.1	62.0	19.3	1.3	0.0
2009	11,420	0.2	5.8	12.5	61.7	18.5	1.2	0.1
2008	11,549	0.3	5.8	12.3	62.4	17.9	1.3	0.1
2007	11,858	0.3	6.0	12.2	63.4	17.1	1.0	0.0
2006	12,457	0.3	5.5	12.3	62.7	17.9	1.2	0.1
2005	11,821	0.3	6.2	12.0	62.7	17.7	1.1	0.1
2004	11,708	0.3	6.2	12.9	62.0	17.3	1.2	0.1
2003	11,814	0.3	6.3	13.3	62.4	16.6	1.0	0.0
2002	11,401	0.3	6.7	14.5	60.7	16.9	1.0	0.0
2001	11,016	0.3	7.2	13.8	61.3	16.3	1.0	0.0
2000	10,381	0.4	7.3	15.0	60.0	16.1	1.2	0.0
1999	9,737	0.4	8.3	14.6	60.2	15.6	0.9	0.0
1998	8,945	0.5	8.3	15.0	59.9	15.4	0.9	0.0
1997	4,132	0.6	9.3	14.8	57.6	16.6	1.1	0.0
1996	7,004	0.6	8.6	14.7	60.0	15.3	1.0	0.0
1995	0	*	*	*	*	*	*	*
1994	7,538	0.6	8.3	14.6	59.4	16.5	0.6	0.0
1993	5,955	0.6	7.6	14.0	61.4	15.8	0.6	0.0
1992	6,190	0.5	7.7	14.0	62.5	14.5	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1991	6,208	0.5	7.3	14.4	63.1	14.0	0.6	0.0
1990	2,259	0.4	6.8	14.5	62.3	15.8	0.2	0.0
1989	2,452	0.7	9.1	14.0	61.7	13.9	0.6	0.0
1988	3,298	0.7	8.5	11.9	64.9	13.5	0.5	0.1
1987	3,790	0.6	8.1	14.6	64.5	11.7	0.4	0.1
1986	2,421	0.6	6.9	13.8	66.2	11.9	0.5	0.2
1985	724	0.6	7.7	12.6	64.9	13.5	0.7	0.0
1984	2,437	0.7	9.0	18.1	60.9	9.7	0.3	1.3
1983	2,492	0.6	9.9	18.5	60.9	9.0	0.2	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2011	9,975	3.7	56.3	9,236	22.2	46.6	1,219	23.9	2,708	19.6
2010	10,533	3.8	55.9	9,871	20.2	47.2	1,307	27.1	3,064	22.9
2009	11,106	3.7	55.2	10,246	20.5	47.5	1,531	27.2	4,177	26.4
2008	11,354	3.8	54.4	10,587	20.8	47.0	1,723	28.1	4,931	26.5
2007	11,646	4.1	52.9	10,900	20.7	46.9	1,631	33.7	5,319	28.6
2006	12,172	4.3	51.7	11,232	20.9	46.8	1,321	28.3	5,413	27.8
2005	11,582	4.3	51.2	10,611	21.3	47.1	1,624	27.8	5,655	29.3
2004	11,510	4.9	50.7	10,472	20.8	48.2	1,509	26.8	5,704	28.4
2003	11,553	4.8	49.4	10,460	25.2	45.1	1,567	25.6	5,606	25.9
2002	11,120	4.9	48.6	10,038	26.3	44.1	1,671	23.9	5,371	24.9
2001	10,810	5.2	47.8	9,881	25.9	42.9	1,607	26.8	4,723	23.4
2000	10,175	5.7	46.3	9,342	25.6	43.2	1,640	26.5	4,499	26.4
1999	9,516	5.9	45.5	8,645	24.3	43.8	1,831	26.1	4,174	23.3
1998	8,687	6.3	44.9	8,006	24.3	44.2	1,595	24.7	3,680	24.0
1997	3,881	5.9	44.0	3,430	22.1	44.1	697	22.2	1,323	19.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1996	6,742	6.7	42.5	6,467	21.4	43.9	1,183	26.3	2,776	21.1
1995	0	*	*	0	*	*	0	*	0	*
1994	7,137	7.7	39.8	6,350	23.5	42.4	1,330	25.0	3,151	19.7
1993	5,716	7.3	39.4	5,314	22.4	42.9	1,065	22.3	2,871	16.1
1992	5,777	7.1	38.4	5,332	25.0	42.2	1,087	22.3	3,165	17.5
1991	5,625	7.5	36.8	5,242	25.0	40.4	1,096	21.8	3,546	16.1
1990	1,901	8.7	35.3	1,785	22.1	42.9	394	19.5	1,416	12.9
1989	2,425	7.6	36.5	0	*	*	622	21.4	588	14.1
1988	3,278	7.9	37.2	0	*	*	982	19.3	258	16.7
1987	3,756	7.7	35.9	0	*	*	1,079	16.6	1,491	16.0
1986	2,368	8.7	34.6	0	*	*	665	15.8	433	8.1
1985	602	10.3	33.6	0	*	*	205	12.7	10	*
1984	2,391	9.7	32.5	0	*	*	705	19.4	1,637	9.8
1983	2,437	10.1	30.0	0	*	*	770	23.9	1,614	12.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2011	9,576	85.7	4.5	10,018	31.2
2010	10,173	87.7	3.3	10,631	28.9
2009	10,566	87.5	2.7	11,153	28.2
2008	10,769	87.0	2.7	11,303	26.5
2007	10,998	88.4	1.9	11,575	27.0
2006	11,298	88.3	1.9	12,196	30.0
2005	10,692	88.5	1.7	11,574	26.7
2004	10,536	88.4	1.7	10,992	29.2
2003	10,574	87.5	2.3	11,587	27.9
2002	10,125	87.4	1.8	11,082	27.4
2001	9,968	87.3	1.7	10,826	25.7
2000	9,412	86.1	1.9	10,177	25.5
1999	8,745	85.7	2.3	8,168	30.0
1998	8,039	84.9	2.1	7,784	28.6
1997	3,205	82.1	1.6	3,919	27.1
1996	5,652	82.4	1.3	6,724	25.9
1995	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
1994	4,353	77.3	2.2	6,056	26.4
1993	2,951	76.8	2.7	4,762	26.0
1992	90	*	*	4,858	20.4
1991	0	*	*	4,646	18.6
1990	0	*	*	1,613	16.4
1989	0	*	*	2,242	21.9
1988	0	*	*	3,140	19.5
1987	0	*	*	3,633	14.4
1986	0	*	*	2,295	12.9
1985	0	*	*	652	18.3
1984	0	*	*	2,320	18.0
1983	0	*	*	2,304	16.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Smoking Indicators

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2011	10,263	31.7	9,558	14.7	2,245	44.0	107	100.0	1,113	100.0
2010	10,802	31.4	10,130	15.5	2,341	41.7	121	100.0	1,477	99.9
2009	11,389	32.7	10,523	15.7	2,478	41.6	207	100.0	2,326	100.0
2008	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
2007	11,750	33.1	10,996	16.3	2,472	41.4	255	100.0	3,320	100.0
2006	12,259	33.9	11,279	16.8	2,640	40.2	261	100.0	3,502	100.0
2005	11,674	33.2	6,175	16.1	1,131	42.3	347	99.7	3,372	100.0
2004	11,564	33.8	2,726	18.0	242	40.5	421	81.9	5,299	66.4
2003	11,665	33.8	1,920	20.0	15	*	1,085	36.4	10,641	32.3
2002	11,224	34.2	2,133	19.4	33	*	1,088	41.2	10,212	32.7
2001	10,913	35.4	2,189	19.8	27	*	920	33.8	10,027	33.4
2000	10,281	35.6	2,257	21.4	13	*	725	40.0	9,472	34.5
1999	9,668	36.4	2,085	21.5	50	*	655	35.0	8,817	34.1
1998	8,867	37.3	2,115	20.9	188	45.2	339	36.0	8,176	34.7
1997	4,015	37.6	1,095	21.0	128	25.8	205	4.4	1,435	35.1
1996	6,737	38.7	2,027	26.7	138	18.8	1	*	0	*
1995	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Smoking Indicators

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
1994	7,041	38.8	1,869	29.2	144	11.1	0	*	0	*
1993	5,535	40.6	1,578	26.5	193	19.7	0	*	0	*
1992	5,592	40.6	1,473	29.1	121	18.2	0	*	0	*
1991	5,154	43.1	1,566	31.7	127	16.5	0	*	0	*
1990	1,044	42.7	570	36.5	26	*	0	*	0	*
1989	0	*	1	*	0	*	0	*	0	*
1988	0	*	0	*	0	*	0	*	0	*
1987	0	*	0	*	0	*	0	*	0	*
1986	0	*	0	*	0	*	0	*	0	*
1985	0	*	0	*	0	*	0	*	0	*
1984	0	*	0	*	0	*	0	*	0	*
1983	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 19C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2011	9,223	7.4	7.0	9,151	9.7	8,232	2.9	9,261	75.6
2010	9,833	7.8	7.4	9,795	10.9	8,696	3.2	9,874	73.5
2009	10,221	7.3	7.4	10,144	10.7	9,033	3.0	10,251	71.2
2008	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
2007	10,795	7.3	7.3	10,611	10.8	9,440	3.1	10,807	69.7
2006	11,118	7.1	7.0	10,937	11.6	9,656	3.0	11,105	69.4
2005	10,520	7.1	7.7	10,377	10.7	9,249	3.1	8,911	99.9
2004	10,370	7.0	7.6	10,249	10.2	9,184	3.2	9,146	64.6
2003	10,397	7.1	7.8	10,238	11.0	9,094	3.2	9,179	62.9
2002	9,996	7.3	8.5	9,765	11.1	8,663	2.9	8,901	60.2
2001	9,778	6.9	8.1	9,653	10.9	8,583	3.1	8,928	58.5
2000	9,202	7.2	8.2	9,056	10.6	8,082	3.2	8,584	58.3
1999	8,499	6.9	8.6	8,299	10.8	7,347	2.9	7,914	53.9
1998	7,823	6.5	8.9	7,378	11.0	6,535	3.0	7,283	50.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1997	3,491	7.9	8.9	3,300	10.9	2,933	3.2	2,368	61.8
1996	6,598	6.5	9.4	6,363	9.9	5,722	2.7	6,345	48.3
1995	0	*	*	0	*	0	*	0	*
1994	6,615	6.9	10.0	5,221	8.8	4,754	3.1	6,237	50.2
1993	5,478	6.5	9.8	4,346	9.7	3,921	2.6	5,184	49.8
1992	5,631	6.2	10.7	4,388	8.9	3,984	2.8	5,079	47.0
1991	5,731	6.5	9.8	4,287	8.3	3,923	3.2	4,714	49.2
1990	2,157	6.8	9.9	1,555	8.4	1,416	3.0	1,364	62.8
1989	2,445	7.9	9.9	2,208	11.7	1,950	4.0	711	100.0
1988	3,286	6.0	10.1	3,073	10.3	2,755	3.5	605	100.0
1987	3,768	5.8	10.9	3,575	12.0	3,138	3.2	712	100.0
1986	2,396	6.5	11.4	2,248	11.0	1,996	3.2	675	100.0
1985	719	6.0	10.6	638	10.2	573	3.1	247	100.0
1984	2,400	6.2	10.4	2,287	11.4	2,021	3.3	754	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1983	2,462	6.4	9.6	2,259	11.4	2,000	3.9	794	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2011	5,636	4.1	55.2	5,215	19.7	50.7	666	21.0	1,611	14.3
	2010	5,918	4.1	56.4	5,526	18.1	50.9	677	21.3	1,810	17.5
	2009	6,123	4.1	54.5	5,646	18.2	51.3	736	21.5	2,103	20.9
	2008	6,456	4.2	54.0	6,013	18.9	51.1	798	25.3	2,581	20.6
	2007	6,633	4.5	53.0	6,190	17.7	51.5	830	28.8	2,767	23.0
	2006	6,974	4.6	50.9	6,421	18.2	51.7	732	22.7	2,836	19.6
	2005	6,667	4.6	50.7	6,126	19.0	51.2	872	24.3	3,057	22.0
	2004	6,721	5.7	49.6	6,130	17.6	53.1	890	21.5	3,043	20.6
	2003	6,879	5.7	47.9	6,206	20.1	50.8	919	21.4	3,070	19.8
	2002	6,763	6.2	47.7	6,092	21.3	49.8	973	20.6	3,093	19.4
Black, Not Hispanic	2011	1,153	4.4	56.3	1,044	25.6	43.7	127	47.2	416	35.8
	2010	1,177	4.6	54.1	1,088	23.7	43.7	146	38.4	454	39.4
	2009	1,214	4.9	56.3	1,080	23.4	47.3	143	51.0	576	46.9
	2008	1,212	4.2	54.2	1,096	22.7	44.5	150	42.7	584	47.6
	2007	1,260	4.8	52.2	1,144	23.5	43.5	169	56.8	673	45.6
	2006	1,315	5.4	50.8	1,150	22.3	43.3	156	52.6	741	44.1
	2005	1,137	4.9	53.0	970	22.8	46.9	126	37.3	704	48.7
	2004	1,113	4.7	51.2	930	20.6	47.2	120	47.5	661	48.6
	2003	1,024	5.3	51.0	878	23.5	44.8	100	49.0	671	42.2
	2002	1,035	4.3	49.8	896	38.4	34.5	176	45.5	687	38.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2011	2,674	1.9	61.2	2,521	25.4	40.7	360	20.3	520	21.9
	2010	2,931	2.4	58.1	2,785	22.5	41.5	407	31.0	626	24.0
	2009	3,294	2.2	58.0	3,092	23.5	41.0	579	28.2	1,331	26.7
	2008	3,182	2.5	57.5	3,017	23.3	40.5	686	26.5	1,549	28.5
	2007	3,259	2.5	55.0	3,107	25.2	39.4	561	34.6	1,663	31.2
	2006	3,383	2.6	55.5	3,200	25.4	38.7	372	28.8	1,629	33.6
	2005	3,257	2.9	53.9	3,058	25.1	40.1	545	29.9	1,657	33.9
	2004	3,131	2.5	55.0	2,925	27.5	39.1	439	32.6	1,746	33.7
	2003	3,106	2.0	54.2	2,892	36.4	34.2	474	27.4	1,568	30.2
	2002	2,845	1.6	51.9	2,614	33.8	34.2	456	22.6	1,353	30.0
American Indian/Alaska Native	2011	142	2.1	64.1	117	22.2	49.6	23	*	34	*
	2010	145	4.1	66.9	133	18.0	57.9	22	*	42	*
	2009	146	3.4	54.1	129	19.4	52.7	19	*	49	*
	2008	178	1.7	57.9	158	19.6	48.1	35	*	67	*
	2007	200	5.5	53.5	182	23.1	45.1	33	*	83	*
	2006	216	3.2	56.0	200	17.5	53.0	24	*	97	*
	2005	217	3.7	53.9	181	19.9	49.7	39	*	96	*
	2004	220	5.0	55.0	191	22.0	50.8	29	*	95	*
	2003	185	3.8	53.0	158	17.7	53.2	30	*	96	*
	2002	192	4.7	56.3	164	20.7	52.4	31	*	79	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2011	256	12.5	26.2	240	33.8	26.7	31	*	106	24.5
	2010	238	10.1	23.5	225	32.4	29.8	35	*	91	*
	2009	214	8.9	26.6	201	22.4	33.8	38	*	87	*
	2008	205	12.2	21.5	194	33.0	32.0	41	*	95	*
	2007	196	12.8	18.4	187	29.4	34.2	27	*	94	*
	2006	211	14.7	18.5	194	30.9	31.4	27	*	80	*
	2005	238	15.1	18.1	219	27.9	29.7	36	*	110	19.1
	2004	244	14.8	19.7	223	23.8	35.9	25	*	117	20.5
	2003	238	17.2	18.9	221	32.1	24.0	32	*	132	23.5
	2002	190	14.2	16.8	182	35.2	28.6	30	*	108	20.4
Multiple Races	2011	113	3.5	54.9	98	*	*	12	*	20	*
	2010	124	4.8	43.5	114	11.4	58.8	20	*	41	*
	2009	115	1.7	49.6	98	*	*	16	*	31	*
	2008	119	4.2	49.6	108	16.7	50.9	13	*	53	*
	2007	98	*	*	90	*	*	11	*	39	*
	2006	73	*	*	67	*	*	10	*	30	*
	2005	64	*	*	57	*	*	6	*	31	*
	2004	29	*	*	29	*	*	3	*	13	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI ⁽³⁾			Weight Gain ⁽⁴⁾			Anemia (Low Hb/Hct) ⁽⁵⁾			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2011	1	*	*	1	*	*	0	*	1	*
	2010	0	*	*	0	*	*	0	*	0	*
	2009	0	*	*	0	*	*	0	*	0	*
	2008	2	*	*	1	*	*	0	*	2	*
	2007	0	*	*	0	*	*	0	*	0	*
	2006	0	*	*	0	*	*	0	*	0	*
	2005	2	*	*	0	*	*	0	*	0	*
	2004	52	*	*	44	*	*	3	*	29	*
	2003	121	3.3	49.6	105	28.6	41.9	12	*	69	*
	2002	95	*	*	90	*	*	5	*	51	*
Total All Race/Ethnic Groups	2011	9,975	3.7	56.3	9,236	22.2	46.6	1,219	23.9	2,708	19.6
	2010	10,533	3.8	55.9	9,871	20.2	47.2	1,307	27.1	3,064	22.9
	2009	11,106	3.7	55.2	10,246	20.5	47.5	1,531	27.2	4,177	26.4
	2008	11,354	3.8	54.4	10,587	20.8	47.0	1,723	28.1	4,931	26.5
	2007	11,646	4.1	52.9	10,900	20.7	46.9	1,631	33.7	5,319	28.6
	2006	12,172	4.3	51.7	11,232	20.9	46.8	1,321	28.3	5,413	27.8
	2005	11,582	4.3	51.2	10,611	21.3	47.1	1,624	27.8	5,655	29.3
	2004	11,510	4.9	50.7	10,472	20.8	48.2	1,509	26.8	5,704	28.4
	2003	11,553	4.8	49.4	10,460	25.2	45.1	1,567	25.6	5,606	25.9
	2002	11,120	4.9	48.6	10,038	26.3	44.1	1,671	23.9	5,371	24.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2011	5,385	87.9	4.3	5,644	32.0
	2010	5,707	89.8	2.8	5,985	30.6
	2009	5,826	90.0	2.7	6,143	30.9
	2008	6,124	89.0	2.7	6,435	29.6
	2007	6,249	89.6	1.9	6,604	29.3
	2006	6,461	90.1	1.5	6,971	31.8
	2005	6,170	90.0	1.6	6,652	29.4
	2004	6,173	89.7	1.6	6,356	31.3
	2003	6,284	89.3	2.1	6,897	29.7
	2002	6,151	89.6	1.6	6,733	29.7
Black, Not Hispanic	2011	1,074	86.9	2.6	1,161	28.5
	2010	1,121	87.8	1.2	1,203	27.2
	2009	1,105	87.8	1.0	1,219	24.4
	2008	1,115	87.4	1.1	1,219	24.5
	2007	1,153	87.7	0.6	1,258	21.5
	2006	1,155	86.5	1.0	1,328	21.5
	2005	975	86.9	0.5	1,147	19.2
	2004	933	88.9	0.2	1,092	21.6
	2003	889	87.5	1.3	1,049	22.1
	2002	907	84.9	0.8	1,042	23.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2011	2,635	81.1	6.0	2,687	31.6
	2010	2,862	83.5	5.4	2,932	27.3
	2009	3,192	83.0	3.4	3,312	25.0
	2008	3,057	83.4	3.2	3,140	21.5
	2007	3,128	86.9	2.4	3,216	24.6
	2006	3,214	85.7	3.0	3,389	29.8
	2005	3,084	86.3	2.0	3,244	23.9
	2004	2,945	85.9	2.3	3,027	27.6
	2003	2,914	83.9	3.1	3,102	25.7
	2002	2,630	83.7	2.4	2,831	23.4
American Indian/Alaska Native	2011	122	81.1	4.9	146	28.1
	2010	137	83.2	2.9	143	22.4
	2009	137	79.6	3.6	147	28.6
	2008	160	80.6	1.3	178	20.2
	2007	188	76.1	1.6	204	25.5
	2006	204	80.9	1.5	223	22.0
	2005	185	81.6	1.6	219	22.4
	2004	189	76.7	5.3	214	25.2
	2003	159	79.9	4.4	181	24.9
	2002	162	79.6	2.5	191	23.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2011	257	84.0	3.1	268	26.1
	2010	226	88.1	1.3	238	19.7
	2009	203	88.7	0.5	215	21.9
	2008	200	89.0	0.5	211	19.9
	2007	189	92.6	1.1	195	23.6
	2006	195	89.7	1.5	212	32.5
	2005	220	89.5	3.2	243	27.2
	2004	223	91.0	0.9	228	29.8
	2003	222	89.6	0.5	235	23.8
	2002	183	89.1	1.1	192	20.8
Multiple Races	2011	102	82.4	4.9	111	26.1
	2010	120	85.8	2.5	130	27.7
	2009	103	87.4	2.9	117	28.2
	2008	111	83.8	4.5	118	28.8
	2007	91	*	*	98	*
	2006	69	*	*	73	*
	2005	58	*	*	66	*
	2004	29	*	*	28	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2011	1	*	*	1	*
	2010	0	*	*	0	*
	2009	0	*	*	0	*
	2008	2	*	*	2	*
	2007	0	*	*	0	*
	2006	0	*	*	0	*
	2005	0	*	*	3	*
	2004	44	*	*	47	*
	2003	106	86.8	0.9	123	39.0
	2002	92	*	*	93	*
Total All Race/Ethnic Groups	2011	9,576	85.7	4.5	10,018	31.2
	2010	10,173	87.7	3.3	10,631	28.9
	2009	10,566	87.5	2.7	11,153	28.2
	2008	10,769	87.0	2.7	11,303	26.5
	2007	10,998	88.4	1.9	11,575	27.0
	2006	11,298	88.3	1.9	12,196	30.0
	2005	10,692	88.5	1.7	11,574	26.7
	2004	10,536	88.4	1.7	10,992	29.2
	2003	10,574	87.5	2.3	11,587	27.9
	2002	10,125	87.4	1.8	11,082	27.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2011	5,776	44.9	5,375	22.5	1,782	40.1	68	*	884	100.0
	2010	6,074	44.7	5,673	23.7	1,876	38.2	85	*	1,124	99.9
	2009	6,281	47.5	5,786	24.4	2,004	38.9	148	100.0	1,782	100.0
	2008	6,550	48.9	6,121	26.2	2,171	36.2	137	100.0	2,536	100.0
	2007	6,692	47.1	6,247	24.9	2,026	38.0	187	100.0	2,551	100.0
	2006	7,020	48.4	6,450	25.7	2,181	36.9	181	100.0	2,722	100.0
	2005	6,722	46.7	3,599	23.8	933	38.5	235	100.0	2,608	100.0
	2004	6,749	47.8	1,707	25.0	194	36.1	293	86.3	3,587	74.6
	2003	6,945	48.2	1,313	26.7	14	*	659	44.3	6,319	42.3
	2002	6,825	48.1	1,395	25.6	28	*	666	48.9	6,199	41.5
Black, Not Hispanic	2011	1,180	22.9	1,072	9.3	185	49.7	16	*	83	*
	2010	1,213	23.8	1,125	10.7	190	45.3	19	*	138	100.0
	2009	1,237	23.8	1,106	10.9	170	42.9	25	*	195	100.0
	2008	1,232	25.7	1,116	13.1	193	36.8	21	*	254	100.0
	2007	1,274	22.4	1,156	10.3	157	48.4	31	*	261	100.0
	2006	1,324	22.5	1,155	10.0	155	47.7	35	*	238	100.0
	2005	1,146	26.4	615	12.2	72	*	53	*	229	100.0
	2004	1,120	23.1	199	14.1	17	*	55	*	437	61.8
	2003	1,040	22.8	133	12.0	0	*	154	32.5	892	27.1
	2002	1,051	21.2	194	14.9	2	*	145	35.9	912	29.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska

Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2011	2,774	8.9	2,631	1.9	180	70.6	9	*	107	100.0
	2010	2,996	8.4	2,852	2.1	182	67.6	10	*	164	100.0
	2009	3,379	8.4	3,189	2.0	197	64.5	19	*	256	100.0
	2008	3,217	8.2	3,054	2.0	188	69.1	15	*	331	100.0
	2007	3,281	8.9	3,125	2.3	201	62.2	23	*	351	100.0
	2006	3,412	8.5	3,208	2.1	191	67.5	32	*	381	100.0
	2005	3,278	8.7	1,689	2.1	89	*	35	*	386	100.0
	2004	3,149	8.3	686	3.4	22	*	49	*	1,015	39.8
	2003	3,135	6.7	383	2.6	0	*	213	13.6	2,941	12.5
	2002	2,870	7.4	458	2.8	1	*	235	21.7	2,654	13.3
American Indian/Alaska Native	2011	148	43.9	121	22.3	46	*	7	*	11	*
	2010	147	47.6	135	11.9	50	*	3	*	24	*
	2009	153	50.3	135	18.5	54	*	8	*	36	*
	2008	179	52.0	158	17.7	57	*	7	*	63	*
	2007	206	45.1	188	14.4	53	*	8	*	80	*
	2006	219	49.3	203	16.7	72	*	7	*	90	*
	2005	220	40.9	112	17.0	25	*	14	*	73	*
	2004	220	52.3	44	*	7	*	14	*	114	70.2
	2003	185	53.5	28	*	1	*	27	*	160	48.8
	2002	190	47.9	35	*	2	*	30	*	168	42.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2011 Pregnancy Nutrition Surveillance (1)
Nebraska

Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2011	269	7.4	256	0.8	12	*	3	*	7	*
	2010	241	8.3	228	2.6	13	*	0	*	9	*
	2009	218	10.1	204	3.4	15	*	2	*	26	*
	2008	211	10.4	200	4.5	15	*	2	*	35	*
	2007	198	10.6	189	3.7	9	*	4	*	40	*
	2006	211	14.2	195	5.1	18	*	3	*	44	*
	2005	241	11.2	136	1.5	5	*	3	*	57	*
	2004	244	10.7	64	*	2	*	7	*	101	56.4
	2003	239	8.8	51	*	0	*	17	*	222	23.9
	2002	192	8.3	42	*	0	*	8	*	186	28.5
Multiple Races	2011	115	48.7	102	17.6	40	*	4	*	21	*
	2010	131	38.9	117	19.7	30	*	4	*	18	*
	2009	121	50.4	103	23.3	38	*	5	*	31	*
	2008	122	47.5	111	19.8	40	*	2	*	50	*
	2007	99	*	91	*	26	*	2	*	37	*
	2006	73	*	68	*	23	*	3	*	27	*
	2005	65	*	24	*	7	*	6	*	19	*
	2004	29	*	11	*	0	*	0	*	15	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2011	1	*	1	*	0	*	0	*	0	*
	2010	0	*	0	*	0	*	0	*	0	*
	2009	0	*	0	*	0	*	0	*	0	*
	2008	2	*	2	*	1	*	0	*	1	*
	2007	0	*	0	*	0	*	0	*	0	*
	2006	0	*	0	*	0	*	0	*	0	*
	2005	2	*	0	*	0	*	1	*	0	*
	2004	53	*	15	*	0	*	3	*	30	*
	2003	121	18.2	12	*	0	*	15	*	107	24.3
	2002	96	*	9	*	0	*	4	*	93	*
Total All Race/Ethnic Groups	2011	10,263	31.7	9,558	14.7	2,245	44.0	107	100.0	1,113	100.0
	2010	10,802	31.4	10,130	15.5	2,341	41.7	121	100.0	1,477	99.9
	2009	11,389	32.7	10,523	15.7	2,478	41.6	207	100.0	2,326	100.0
	2008	11,513	34.4	10,762	17.4	2,665	39.1	184	100.0	3,270	100.0
	2007	11,750	33.1	10,996	16.3	2,472	41.4	255	100.0	3,320	100.0
	2006	12,259	33.9	11,279	16.8	2,640	40.2	261	100.0	3,502	100.0
	2005	11,674	33.2	6,175	16.1	1,131	42.3	347	99.7	3,372	100.0
	2004	11,564	33.8	2,726	18.0	242	40.5	421	81.9	5,299	66.4
	2003	11,665	33.8	1,920	20.0	15	*	1,085	36.4	10,641	32.3
	2002	11,224	34.2	2,133	19.4	33	*	1,088	41.2	10,212	32.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2011	5,209	6.9	7.4	5,169	9.8	4,647	2.4	5,226	73.7
	2010	5,519	8.0	8.0	5,501	10.6	4,899	3.2	5,540	70.7
	2009	5,635	6.9	7.8	5,585	10.4	4,998	2.6	5,650	67.6
	2008	5,942	7.0	7.3	5,907	10.2	5,293	2.8	5,957	66.7
	2007	6,125	7.0	7.7	6,037	10.7	5,382	2.8	6,133	66.2
	2006	6,358	6.8	7.4	6,252	11.3	5,537	2.7	6,353	65.8
	2005	6,076	6.9	8.2	5,993	10.5	5,352	2.8	5,129	99.9
	2004	6,074	6.8	8.2	5,989	10.6	5,353	2.7	5,330	59.9
	2003	6,169	7.0	8.4	6,079	11.4	5,376	3.0	5,456	57.7
	2002	6,051	7.3	8.7	5,918	11.5	5,219	2.8	5,404	55.5
Black, Not Hispanic	2011	1,042	11.7	4.3	1,037	11.0	919	5.0	1,047	70.1
	2010	1,093	12.4	3.8	1,094	13.5	944	5.4	1,097	67.7
	2009	1,074	12.5	4.1	1,069	12.5	931	7.2	1,079	64.0
	2008	1,085	12.4	4.7	1,079	12.8	939	5.6	1,088	59.6
	2007	1,140	12.3	3.8	1,123	12.6	977	5.6	1,146	60.1
	2006	1,156	11.4	4.2	1,142	14.2	980	5.4	1,151	61.5
	2005	964	11.1	4.9	952	13.2	825	5.3	804	100.0
	2004	929	10.3	5.3	932	10.3	829	5.2	890	53.8
	2003	881	9.5	4.0	873	10.9	775	4.1	842	56.5
	2002	901	11.1	5.3	881	12.9	767	4.2	871	49.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2011	2,508	6.5	7.8	2,483	9.3	2,242	2.7	2,522	82.4
	2010	2,758	5.9	7.6	2,739	10.3	2,448	2.3	2,771	81.5
	2009	3,081	6.0	7.8	3,063	10.6	2,727	2.1	3,090	80.4
	2008	2,951	6.9	6.6	2,922	10.3	2,614	3.0	2,957	81.7
	2007	3,068	6.0	7.4	2,998	10.5	2,680	2.6	3,066	81.5
	2006	3,142	6.0	7.3	3,084	11.2	2,737	2.8	3,140	79.8
	2005	3,021	6.3	7.3	2,975	10.5	2,657	2.9	2,588	100.0
	2004	2,891	6.4	7.4	2,856	9.4	2,583	3.4	2,494	79.1
	2003	2,869	6.3	7.7	2,816	9.9	2,533	3.2	2,459	77.8
	2002	2,610	6.0	8.9	2,550	9.7	2,300	2.8	2,243	76.4
American Indian/Alaska Native	2011	118	4.2	10.2	116	6.9	108	3.7	118	65.3
	2010	129	4.7	8.5	129	10.9	114	1.8	131	60.3
	2009	134	7.5	10.4	131	11.5	116	2.6	134	70.1
	2008	155	6.5	12.3	156	10.9	138	3.6	156	69.9
	2007	184	6.0	14.7	179	11.7	158	1.9	183	63.9
	2006	202	6.9	8.4	201	12.9	174	4.0	201	64.7
	2005	184	4.3	15.2	183	11.5	162	0.6	148	100.0
	2004	190	8.4	12.6	186	12.9	162	2.5	171	54.4
	2003	156	5.8	12.2	152	14.5	130	1.5	138	51.4
	2002	158	3.2	15.2	149	9.4	135	0.7	146	54.1

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
Asian/Pacific Islander	2011	244	10.2	2.9	245	9.0	221	5.9	246	76.8
	2010	219	6.4	6.4	217	11.1	193	3.1	220	77.3
	2009	200	8.0	4.0	200	11.0	177	4.5	201	70.6
	2008	196	7.7	2.0	194	10.3	174	2.9	196	71.9
	2007	188	9.0	6.9	186	10.2	166	3.6	189	58.7
	2006	192	9.4	5.2	190	12.6	166	2.4	192	70.3
	2005	217	7.4	5.1	216	7.9	199	4.5	191	100.0
	2004	218	6.4	3.7	218	10.6	194	3.6	197	65.0
	2003	213	8.5	4.7	213	12.2	186	5.4	189	48.1
	2002	184	9.8	4.3	179	8.4	163	5.5	164	47.6
Multiple Races	2011	101	5.9	4.0	100	6.0	94	*	101	69.3
	2010	115	5.2	6.1	115	14.8	98	*	115	75.7
	2009	97	*	*	96	*	84	*	97	*
	2008	109	5.5	5.5	106	10.4	95	*	109	58.7
	2007	90	*	*	88	*	77	*	90	*
	2006	68	*	*	68	*	62	*	68	*
	2005	58	*	*	58	*	54	*	51	*
	2004	29	*	*	28	*	27	*	27	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Nebraska

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2011	1	*	*	1	*	1	*	1	*
	2010	0	*	*	0	*	0	*	0	*
	2009	0	*	*	0	*	0	*	0	*
	2008	2	*	*	2	*	2	*	2	*
	2007	0	*	*	0	*	0	*	0	*
	2006	0	*	*	0	*	0	*	0	*
	2005	0	*	*	0	*	0	*	0	*
	2004	39	*	*	40	*	36	*	37	*
	2003	109	9.2	9.2	105	9.5	94	*	95	*
	2002	92	*	*	88	*	79	*	73	*
Total All Race/Ethnic Groups	2011	9,223	7.4	7.0	9,151	9.7	8,232	2.9	9,261	75.6
	2010	9,833	7.8	7.4	9,795	10.9	8,696	3.2	9,874	73.5
	2009	10,221	7.3	7.4	10,144	10.7	9,033	3.0	10,251	71.2
	2008	10,440	7.5	6.8	10,366	10.5	9,255	3.2	10,465	70.3
	2007	10,795	7.3	7.3	10,611	10.8	9,440	3.1	10,807	69.7
	2006	11,118	7.1	7.0	10,937	11.6	9,656	3.0	11,105	69.4
	2005	10,520	7.1	7.7	10,377	10.7	9,249	3.1	8,911	99.9
	2004	10,370	7.0	7.6	10,249	10.2	9,184	3.2	9,146	64.6
	2003	10,397	7.1	7.8	10,238	11.0	9,094	3.2	9,179	62.9
	2002	9,996	7.3	8.5	9,765	11.1	8,663	2.9	8,901	60.2

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
(3) <2500 g.
(4) >4000 g.
(5) <37 weeks gestation.
(6) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

