
CDC NEVDRS - 2022


NEVDRS NEBRASKA VIOLENT DEATH **REPORTING SYSTEM**

Summary of Suicide Deaths in Behavioral Health Region 6 in Nebraska

120 Total Deaths (14.0 per 100,000 Population)




In 2022, the majority of suicide victims were white males aged 40-49 with a high school education in Nebraska Behavioral Health Region 6. Asian/Pacific Islander and Black Non-Hispanic males were disproportionately affected by suicide. In 2021, the most impacted age group was 30-39 years old, which was the main difference between the two years.




In 2022, around 30% of suicides in Nebraska Behavioral Health Region 6 occurred in the fourth guarter, with February and November recording the highest numbers, accounting for 23.4% of the total. The majority of the victims were white males, with an average age of 40 and at least a high school education during the final quarter. In 2021, most suicide deaths also occurred in the fourth quarter, though there was a slightly higher proportion of deaths in the first quarter compared to the second and third quarters.




Suicide Death Rate by Census Tracts





In 2022, firearms were the most common method of suicide among males in Nebraska Behavioral Health Region 6, with the average age of firearm-related deaths at 44. Strangulation, occurring at an average age of 41, was less common. In 2021, firearm deaths were about 20% lower, while strangulation was 11% more frequent, with average ages remaining similar.



The darker blue areas on this map represent the highest rate of suicide deaths in Nebraska Behavioral Health Region 6 in 2022.

30 miles

Other Decedent Characteristics

NFBRASKA

DEPT. OF HEALTH AND HUMAN SERVICES

Statistics are b	ased on 118 cases.	Categories are not mutually exclusive.
	79.7%	of total suicide deaths took place in a residence.
	11.9 %	of total suicide victims experienced homelessness or housing instability.
	36.4%	of total suicide victims had previous mental health problems.
	26.3%	of total suicide victims were diagnosed with depression.
	7.6%	of total suicide victims were diagnosed with anxiety disorder.
	28.8%	of total suicide victims had history of mental health treatment.
	24.6 %	of total suicide victims had current mental health treatment.
	32.2%	of total suicide victims were perceived by others to be depressed at the time of death.
	33.1%	of total suicide victims had a history of suicidal ideation.
	21.2 %	of total suicide victims had a history of suicide attempt.
	27.1%	of total suicide victims had recently disclosed their suicidal intent.
	10.2%	of total suicide victims had disclosed their suicidal intent to their friend.
	33.1 %	of total suicide victims left suicide notes.
	28.0%	of total suicide victims had a history of alcohol abuse.
	17.8%	of total suicide victims had a history of substance abuse other than alcohol.
	28.0%	of total suicide victims had a history of physical health problems.
The reduced death of	7.6%	of total suicide victims had criminal or legal problems.
	29.7 %	of total suicide victims had intimate partner problems.
	12.7%	of total suicide victims had family or other relationship problems.
	15.3%	of total suicide victims had known recent argument.
	44.9%	of the suicide deaths were among never married individuals.
	23.7%	of the suicide deaths were among divorced individuals.

Due to small numbers, rates may be unstable.

The reduced death count in this section is due to incomplete law enforcement or coroner reports.