

## DRUGS

Anything that gets you high can harm your baby. Stay away from marijuana, crack, cocaine, heroin, ecstasy, meth, and other street drugs.

Using street drugs, while pregnant increases the risk that your baby could:

- Be born too small, too early, or stillborn
- Be born addicted to drugs
- Have a birth defect, deformities, or trouble breathing
- Have learning, behavioral, or other health problems throughout life

Get Help. Stopping drugs suddenly without a doctor's help can cause serious withdrawal and health problems for you and your baby.



### What to do now:

- Protect your baby. Never use drugs.
- Ask for help if you're using drugs. Even if you had trouble quitting before, try again now for your baby's health.
- Only take prescription drugs that your doctor has approved for you during your pregnancy and/or while breastfeeding.

### What to do when the baby comes:

- Stay drug-free. Your baby depends on you for love and care.
- Remember drugs pass into your breast milk. A healthy lifestyle means better milk for your baby.

**If you, or someone you know, would like help to quit, try these resources:**



For help quitting smoking, call 1-800-QUIT NOW (1-800-784-8669) or scan the QR Code to visit <https://ne.quitlogix.org/en-us/>

For help quitting drinking or drugs, call 1-800-NCA-CALL (1-800-622-2255)

*This institution is an equal opportunity provider*

Source: USDA, Food and Nutrition Services FNS-489

# KEEP YOURSELF AND YOUR BABY SAFE

EXPOSING THE DANGERS OF SMOKING, DRINKING AND TAKING DRUGS DURING PREGNANCY AND WHILE BREASTFEEDING

## SMOKING

Tobacco contains harmful chemicals that can hurt your baby.

Not smoking gives your baby a healthy start! Smoking while pregnant increases the risk that your baby could:

- Be born too small or too early
- Be stillborn or die during infancy
- Have breathing problems or trouble sleeping
- Have behavioral problems later in life

Exposure to secondhand smoke increases your baby's risk of asthma, ear infections, allergies, pneumonia, and SIDS (sudden infant death syndrome).



### What to do now:

- If you use tobacco products, QUIT TODAY—it's never too late. Ask for help; you don't have to do it alone.
- Even if you had trouble quitting before, try again for the health of your baby.

### What to do when the baby comes:

- Keep your baby away from smoke, and don't let anyone else smoke around your baby.
- Breastfeed your baby, and avoid cigarettes. Smoking can lower your milk supply and make it harder for your baby to gain weight.

## ALCOHOL

Beer, wine, wine coolers, and mixed drinks are all harmful to your baby.

### **No amount of alcohol is safe while you are pregnant.**

Even if you drank before you knew you were pregnant, stop now for the sake of your baby's health. Drinking while pregnant can lead to fetal alcohol spectrum disorders, including:

- Birth defects such as a deformed face
- Problems seeing or hearing
- Trouble learning
- Mental and behavioral issues

Drinking while pregnant also increases the risk that your baby could be miscarried, born too early, or stillborn.



### **What to do now:**

- Do not drink while you are pregnant.
- If quitting is hard, ask for help. There are ways to make it easier to stop drinking.

### **What to do when the baby comes:**

- Be very cautious about drinking alcohol, if you choose to drink at all.
- You may consume a single alcoholic drink if your baby's breastfeeding routine is well established - no earlier than at 3 months old. Then wait at least 4 hours after having a drink before breastfeeding. Or express breast milk before drinking to feed your baby later.
- Remember, your baby depends on you. Be responsible. Stay sober.



## MEDICATION

Ask your doctor before taking any medicines. Medicines you used before you got pregnant may not be safe to take now. Even those sold at a drugstore or grocery store can hurt your baby.

Taking certain medicines while pregnant increases the risk that your baby could:

- Be born too small, too early, or stillborn
- Have high-pitched crying and be hard to comfort
- Have a birth defect, developmental problems, or seizures.

Misusing prescription drugs can also harm your baby. This includes taking your own prescription in a way not intended by your doctor and taking a prescription drug prescribed for someone else.

**Only take medicines your doctor says are safe.**

### **What to do now:**

- Use medicines wisely. Tell your doctor about all medicines or supplements you take to make sure it's safe to keep taking them.
- If you get sick, ask your doctor which medicines are safe to take.

### **What to do when the baby comes:**

- Check with your doctor before taking medicines while you are breastfeeding. Medicines can get into your baby through your breast milk.

