


CDC NeVDRS - 2020 and 2021

Summary of Male Suicide Deaths in Nebraska


461 Total Deaths (23.8 per 100,000 Population)






AI/AN, NH: American Indian/Alaskan Native, non-Hispanic. A/PA, NH: Asian or Pacific Islander, non-Hispanic. * Suppressed due to small numbers.


In 2020 and 2021, among male suicide victims, white males with a high school education and between the age of 30-39 years old most frequently died from suicide. American Indian/Alaska Native, Non-Hispanic and Black, Non-Hispanic males were disproportionately impacted by suicide in Nebraska.



In 2020 and 2021, about 28.0% of male suicides in Nebraska occurred in the last quarter. The month of November registered the highest number of suicides, comprising 10.7% of the total. The predominant demographic among individuals who lost their lives to suicide during these months were white males, with an average age of 43 and holding a high school degree.



Suicide Death Rate by Census Tracts



Other Decedent Characteristics		
Statistics are based on 403 cases. Categories are not mutually exclusive.		
	72.2%	of total suicide deaths took place in a residence.
	11.2%	of total suicide victims experienced homelessness or housing instability.
	38.0%	of total suicide victims had previous mental health problems.
	28.0%	of total suicide victims were diagnosed with depression.
	7.9%	of total suicide victims were diagnosed with anxiety disorder.
	23.1%	of total suicide victims had history of mental health treatment.
	14.1%	of total suicide victims had current mental health treatment.
	60.5%	of total suicide victims were perceived by others to be depressed at the time of death.
	29.5%	of total suicide victims had a history of suicidal ideation.
	12.4%	of total suicide victims had a history of suicide attempt.
	26.3%	of total suicide victims had recently disclosed their suicidal intent.
	12.2%	of total suicide victims had disclosed their suicidal intent to their intimate partner.
	30.3%	of total suicide victims left suicide notes.
	25.9%	of total suicide victims had a history of alcohol abuse.
	20.3%	of total suicide victims had a history of substance abuse other than alcohol.
	22.8%	of total suicide victims had a history of physical health problems.
	12.7%	of total suicide victims had criminal or legal problems.



27.5%

of total suicide victims had family or other 8.7% relationship problems.

problems.

of total suicide victims had intimate partner

of total suicide victims had known recent 17.4% argument.

of the suicide deaths were among never 44.9% married individuals.

of the suicide deaths were among married 27.0% individuals.