


81 Total Deaths (2.1 per 100,000 Population)

Demographic Information




In 2020 and 2021, among veterans, white males with a high school education or less and older than 70 years old most frequently died from suicide. American Indian/Alaska Native and Hispanic males were disproportionately impacted by suicide in Nebraska.


Cause of Death



Suicide Deaths by Month



Suicide Death Rate by Census Tracts



Other Decedent Characteristics

- Statistics are based on 74 cases. Categories are not mutually exclusive.
- 77.0% of total suicide deaths took place in a residence.
 - 18.9% of total suicide victims experienced homelessness or housing instability.
 - 37.8% of total suicide victims had previous mental health problems.
 - 25.7% of total suicide victims were diagnosed with depression.
 - 8.1% of total suicide victims were diagnosed with anxiety disorder.
 - 18.9% of total suicide victims had history of mental health treatment.
 - 13.5% of total suicide victims had current mental health treatment.
 - 64.9% of total suicide victims were perceived by others to be depressed at the time of death.
 - 20.3% of total suicide victims had a history of suicidal ideation.
 - 20.3% of total suicide victims had a history of suicide attempt.
 - 20.3% of total suicide victims had recently disclosed their suicidal intent.
 - 8.1% of total suicide victims had disclosed their suicidal intent to their family member.
 - 28.4% of total suicide victims left suicide notes.
 - 27.0% of total suicide victims had a history of alcohol abuse.
 - 9.5% of total suicide victims had a history of substance abuse other than alcohol.
 - 33.8% of total suicide victims had a history of physical health problems.
 - 8.1% of total suicide victims had criminal or legal problems.
 - 20.3% of total suicide victims had intimate partner problems.
 - 8.1% of total suicide victims had family or other relationship problems.
 - 16.2% of total suicide victims had known recent argument.
 - 40.5% of the suicide deaths were among married individuals.
 - 27.0% of the suicide deaths were among divorced individuals.