


CDC NeVDRS - 2020 and 2021

Summary of Working Age Population Suicide Deaths in Nebraska


DEPT. OF HEALTH AND HUMAN SERVICES

441 Total Deaths (11.3 per 100,000 Population)




AI/AN, NH: American Indian/Alaskan Native, non-Hispanic. A/PA, NH: Asian or Pacific Islander, non-Hispanic.


In 2020 and 2021, among working age individuals, white males with a high school education and between the age of 30-39 years old most frequently died from suicide. American Indian/Alaska Native and Hispanic males were disproportionately impacted by suicide in



In 2020 and 2021, approximately 28.4% of working age population suicides in Nebraska took place in the last quarter. The month of October recorded the highest number of suicides, accounting for 11.4% of the total. The majority of individuals who lost their lives to suicide during these months were white males, with an average age of 40 and possessing a high school degree. Notably, Asian/Pacific Islander females and Black males in Nebraska were disproportionately impacted by suicide.



Suicide Death Rate by Census Tracts





C	ther Dece	dent Characteristics
Statistics	are based on 391 cases.	Categories are not mutually exclusive.
	72.6%	of total suicide deaths took place in a residence.
	12.8%	of total suicide victims experienced homelessness or housing instability.
	44.2%	of total suicide victims had previous mental health problems.
	32.7%	of total suicide victims were diagnosed with depression.
	9.2%	of total suicide victims were diagnosed with bipolar disorder or anxiety disorder.
	29.2%	of total suicide victims had history of mental health treatment.
	18.2%	of total suicide victims had current mental health treatment.
	59.6%	of total suicide victims were perceived by others to be depressed at the time of death.
	33.5%	of total suicide victims had a history of suicidal ideation.
	17.4%	of total suicide victims had a history of suicide attempt.
	26.9%	of total suicide victims had recently disclosed their suicidal intent.
	26.9%	of total suicide victims had disclosed their suicidal intent to their intimate partner.
	32.2%	of total suicide victims left suicide notes.
	27.6%	of total suicide victims had a history of alcohol abuse.
	27.1%	of total suicide victims had a history of substance abuse other than alcohol.
	15.9%	of total suicide victims had a history of physical health problems.
	13.3%	of total suicide victims had criminal or legal problems.
		of total acciding violing bod between a section

31.2%

9.2%

44.2%

27.6%

problems.

argument.

individuals.

relationship problems.

married individuals.

of total suicide victims had intimate partner

of total suicide victims had family or other

of total suicide victims had known recent

of the suicide deaths were among never

of the suicide deaths were among married