Attachment to the Facility Worksheet for the Live Birth Certificate for Multiple Births

This attachment is to be completed when at least two infants in a multiple pregnancy are born alive. Complete a full worksheet for the first-born infant and an attachment for each additional live-born infant. A "Facility Worksheet for the Report of Fetal Death" should be completed for any fetal loss in this pregnancy reportable under State reporting requirements. For "Delayed Interval Births," that is, births in a multiple pregnancy delivered at least 24 hours apart, a full worksheet, not an attachment should be completed.

Mother's name:
Mother's medical record #
Child's name:
Child's medical record #
Mother's Pregnancy History (Sources: Prenatal care records, mother's medical records)
Number of previous live births now living (For this multiple delivery, include live born infants born before this infant in the delivery who are still living, in addition to infants from prior pregnancies. If none enter "0"):
Number of previous live births now dead (For this multiple delivery, include live born infants born before this infant in the delivery who are now dead, in addition to infants from prior pregnancies. If none enter "0"):
Date of last live birth: Month Year
Total number of other pregnancies (For this multiple delivery, include all fetal losses of any gestational age delivered before this infant in the pregnancy, in addition to losses from prior pregnancies. If none enter "0"):
Date of last other pregnancy outcome (Date when last pregnancy which did not result in a live birth ended): Month Year
<u>Labor and Delivery</u> (Sources: Labor and delivery records, mother's medical records)
Date of birth: Month Day Year
Time of birth (12-hour clock):Time (Circle one) a.m. p.m. noon midnight
Order delivered in the pregnancy (Specify born 2nd, 3rd, 4th, 5th, 6th, 7th, etc.) (Include all live births and fetal losses resulting from this pregnancy):
Characteristics of labor and delivery (Check all that apply to this infant): Induction of labor - (Initiation of uterine contractions by medical and/or surgical means for the purpose of delivery before the spontaneous onset of labor.) Augmentation of labor - (Stimulation of uterine contractions by drug or manipulative technique with the intent to reduce the time to delivery.) Non-vertex presentation - (Includes any non-vertex fetal presentation, e.g., breech, shoulder, brow, face presentations, and transverse lie in the active phase of labor or at delivery other than vertex.) Steroids (glucocorticoids) for fetal lung maturation received by the mother prior to delivery - (Includes betamethasone, dexamethasone, or hydrocortisone specifically given to accelerate fetal lung maturation in anticipation of preterm delivery. Excludes steroid medication given to the mother as an anti-inflammatory treatment.) Antibiotics received by the mother during labor - (Includes antibacterial medications given systemically)
Antibiotics received by the mother during labor - (Includes antibacterial medications given systemically (intravenous or intramuscular) to the mother in the interval between the onset of labor and the actual delivery: Ampicillin, Penicillin, Clindamycin, Erythromycin, Gentamicin, Cefataxime, Ceftriaxone, etc.)

intensive care.) Unplanned operating room procedure following of	•	
Includes anticipated but not definitively planned Admission to intensive care unit - (Any admission	hysterectomy.)	
Ruptured uterus - (Tearing of the uterine wall.)Unplanned hysterectomy - (Surgical removal of the uterine wall)	he literus that was not planned prior to the	admission
vaginal mucosa, perineal body and anal sphincter rectal mucosa.)	r. 4° laceration is all of the above with ext	ension through the
Third or fourth degree perineal laceration - (3° la		
Maternal transfusion - (Includes infusion of whole delivery.)	le blood or packed red blood cells associat	ed with labor and
Maternal morbidity (Serious complications experience (Check all that apply to <u>this</u> infant):	ced by the mother associated with labor ar	nd delivery)
with plans for a vaginal delivery.) (Circle One) Yes No	or unempiea: - (Labor was anowed, augh	icinca or induced
to the fetal head.) ——Cesarean - (Extraction of the fetus, placenta and and uterine walls.) If cesarean, was a trial of lab		
Vaginal/Vacuum - (Delivery of the fetal head thro	ough the vagina by application of a vacuur	n cup or ventouse
<i>Vaginal/Forceps</i> - (Delivery of the fetal head throfetal head.)		al forceps to the
<i>Vaginal/Spontaneous</i> - (Delivery of the entire fett without manual assistance from the delivery atter	ndant.)	
Other - (Any other presentation not listed above) D. Final route and method of delivery (Check one):		
Breech - (Presenting part of the fetus listed as bre		
C. Fetal presentation at birth (Check one): Cephalic - (Presenting part of the fetus listed as v	ertex, occiput anterior (OA), occiput posto	erior (OP))
If Yes, Was it successful? (Circle One) Yes	No	
B. Was delivery with vacuum extraction attempted? - (attempt at vaginal delivery.) (Circle One) Yes	(Ventouse or vacuum cup was applied to the No	he fetal head in an
delivery.) (Circle One) Yes No If Yes, Was it successful? (Circle One) Yes	No	
(Complete A, B, C, and D): A. Was delivery with forceps attempted? - (Obstetric in the linear of the Complete	forceps was applied to the fetal head in an	attempt at vaginal
Method of Delivery (The physical process by which t	he complete delivery of the infant was effe	ected)
None of the above		
of the pain of labor, i.e., delivery of the agent int limited to the lower body.)	o a limited space with the distribution of t	he analgesic effect
Epidural or spinal anesthesia during labor - (Ad		
Operative delivery – operative intervention to sh cesarean delivery.)		
the mother, amnioinfusion, support of maternal be Further fetal assessment includes any of the following		
following - maternal position change, oxygen ad	ministration to the mother, intravenous flu	ids administered to
Fetal intolerance of labor was such that one or m measures, further fetal assessment, or operative		
change of an otherwise clear fluid.)		
Moderate/heavy meconium staining of the amnion fetal bowel contents during labor and/or at deliver		
temperature at or above 38°C (100.4°F).	madility, leukocytosis and letai tacifycard	ia. Airy maternai
diagnosis of chorioamnionitis during labor made the following: fever, uterine tenderness and/or in	by the delivery attendant. Usually include	es more than one of
Clinical chorioamnionitis diagnosed during labor	r or maternal temperature $>=38^{\circ}$ C (100.	4° F) - (Clinical

Newborn

(Sources: Labor and delivery records, Newborn's medical records, mother's medical records)

Birthweight (grams p	referred):		grams or	pounds and	ounces
	final estimat	te of gestation b	ased on all perinat	al factors and assessments, beriod and the date of birth.)	out not the
Sex (Circle One)	Male	Female	Not yet determi	ned	
Apgar score (A system birth):	natic measu	re for evaluating	g the physical cond	dition of the infant at specific	e intervals at
Score at 5 minutes		Score at 10 m	inutes (If 5 minute	e score is less than 6)	
Check all that apply):	ion required ask or bag a oscopy for a ion required my method for a new surfactant results as both artification of the surfactant of the s	d immediately for and endotracheal aspiration of medifor more than for > 6 hours. Incoming a facility whorn.) eplacement there deficiency due to deficiency due to decial and extracted whom for suspections, etc.) give ic dysfunction - ic dysfunction is a cencephalopath de symptoms associated fracture(s) intervention) - ic yery. Includes an intervention is and/or hypoten leen, or adrenal	allowing delivery - tube within the first conium.) six hours - (Infant cludes conventionally or unit staffed an apy - (Endotrachea preterm birth or ped natural surfactate preterm birth or ped natural surfactates extensive any invisevere alteration or period with CNS, peripheral nerve (Defined as presently bony fracture or extravasation within and/or extremity of sion. Solid organ I hematoma.)	psis - (Any antibacterial drug ntravenous or intramuscular) voluntary repetitive, convulsi of alertness such as obtundat rgy or hypotonia in the absen congenital anomalies.) injury, and/or soft tissue/sol at immediately following delate weakness or loss of sensation to the hemorrhage requiring evaluation the scalp) hemorrhage, gial ecchymosis accompanied by themorrhage includes subcaps	s for any duration h. Excludes oxygen n (breathing ntinuous positive nuous mechanical tive suspension for n respiratory g (e.g., penicillin,).) tive movement or tion, stupor, or nce of other lid organ livery or on, but excludes nation and/or evidence of sular hematoma of
(Check all that apply): Anencephaly - (2) absent brain. AlMeningomyeloc. a bony defect of Meningocele (h Both open and of midline bony spCyanotic congection in the congection of the congenital diagrams.	Partial or co so includes ele/Spina big f spine closu erniation of closed (coverinal defect position of the datresia, true thragmatic in the solution of the s	mplete absence infants with cra fida - (Spina bifure. Meningomy meninges without red with skin) le without protrusidisease - (Conge the great arteries incus arteriosus, thernia - (Defect	of the brain and sk niorachischisis (an ida is herniation of elocele is herniation out spinal cord tiss esions should be in on of the spinal co- enital heart defects is (vessels), tetratol total/partial anom	corn diagnosed prenatally or a cull. Also called anencephalus hencephaly with a contiguous of the meninges and/or spinal con of meninges and spinal con ue) should also be included included. Do not include Spin ord or meninges).) which cause cyanosis. Inclu- ogy of Fallot, pulmonary or alous pulmonary venous retu- of the diaphragm allowing he	as, acrania, or s spine defect). cord tissue through ord tissue. in this category. It is bifida occulta (a des but is not pulmonic valvular urn with or without
obstruction.)	ohragmatic l	hernia - (Defect			

Medical Record Numbers—Mother's: _____ Child's: _____

Page 3 of 4

Omphalocele - (A defect in the anterior abdominal wall, accompanied by herniation of some abdominal	
organs through a widened umbilical ring into the umbilical stalk. The defect is covered by a membrane	
(different from gastroschisis, see below), although this sac may rupture. Also called exomphalos. Do not	
include umbilical hernia (completely covered by skin) in this category.)	
Gastroschisis - (An abnormality of the anterior abdominal wall, lateral to the umbilicus, resulting in	
herniation of the abdominal contents directly into the amniotic cavity. Differentiated from omphalocele b	y
the location of the defect and absence of a protective membrane.)	
Limb reduction defect (excluding congenital amputation and dwarfing syndromes) - (Complete or partial	
absence of a portion of an extremity associated with failure to develop.)	
Cleft Lip with or without Cleft Palate - (Incomplete closure of the lip. May be unilateral, bilateral or	
median.)	
Cleft Palate alone - (Incomplete fusion of the palatal shelves. May be limited to the soft palate or may	
extend into the hard palate. Cleft palate in the presence of cleft lip should be included in the "Cleft Lip w	ith
or without Cleft Palate" category above.)	
Down Syndrome - (Trisomy 21) (Circle One) Karyotype confirmed Karyotype pending	
Suspected chromosomal disorder - (Includes any constellation of congenital malformations resulting from	n or
compatible with known syndromes caused by detectable defects in chromosome structure.)	
(Circle One) Karyotype confirmed Karyotype pending	
Hypospadias - (Incomplete closure of the male urethra resulting in the urethral meatus opening on the ven	ıtral
surface of the penis. Includes first degree - on the glans ventral to the tip, second degree - in the coronal	
sulcus, and third degree - on the penile shaft.)	
None of the above	
Was infant transferred within 24 hours of delivery? (Circle "yes" if the infant was transferred from this facility to another within 24 hours of delivery. If transferred more than once, enter name of first facility to which the integral was transferred.) (Circle One) Yes No If Yes, name and location of the facility infant transferred to:	
Infant living at time of report? (Infant is living at the time this birth certificate is being completed. Answer "Y	es"
if the infant has already been discharged to home care.)	
YesNoInfant transferred, status unknown	
Is infant being breast fed at discharge? (Circle One) Yes No	
Is infant going to be adopted? (Circle One) Yes No Don't know	
If the Acknowledgement of Paternity form has been signed, provide the date signed (if the date signed by t	
in the field wiedgement of i decimely form has been signed, provide the date signed in the date signed by	he
	he
mother and father are different, provide the latest date signed): Month DayYear	he
	he
	he
	he